

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 February 2002 (21.02.2002)

PCT

(10) International Publication Number
WO 02/14317 A2

- (51) International Patent Classification⁷: C07D 471/04, 519/00, A61K 31/437, A61P 29/00, 11/06 // (C07D 471/04, 231:00, 221:00) (C07D 519/00, 471:00, 471:00)
- (21) International Application Number: PCT/US01/25180
- (22) International Filing Date: 10 August 2001 (10.08.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/225,178 14 August 2000 (14.08.2000) US
09/927,188 10 August 2001 (10.08.2001) US
- (71) Applicant: **ORTHO MCNEIL PHARMACEUTICAL, INC.** [US/US]; U.S. Route 202, P.O. Box 300, Raritan, NJ 08869-0602 (US).
- (72) Inventors: **CAI, Hui**; 11213 Corte Belleza, San Diego, CA 92130 (US). **EDWARDS, James, P.**; 8723 Hesby Court, San Diego, CA 92129 (US). **MEDUNA, Steven, P.**; 9975 Paseo Montril, San Diego, CA 92129 (US). **PIO, Barbara, A.**; 3898 Ashford Street, San Diego, CA 92111 (US). **WEI, Jianmei**; 12516 Senda Panacea, San Diego, CA 92129 (US).
- (74) Agents: **JOHNSON, Philip, S. et al.**; One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SUBSTITUTED PYRAZOLES

(57) Abstract: Substituted pyrazoles, methods of manufacturing them, compositions containing them, and methods of using them to treat, for example, autoimmune diseases mediated by cathepsin S are described.



WO 02/14317 A2

SUBSTITUTED PYRAZOLES

Field of the Invention

5 This invention relates to a series of substituted pyrazoles, pharmaceutical compositions containing these compounds, and intermediates used in their manufacture, and methods of using them.

Background of the Invention

10

Cathepsin S (EC 3.4.22.27) is a cysteine protease of the papain family found primarily in lysosomes (Bromme, D.; McGrath, M. E. High Level Expression and Crystallization of Recombinant Human Cathepsin S. *Protein Science* **1996**, *5*, 789-791).

15

The role of cathepsin S in the immune response is anticipated by its tissue distribution: cathepsin S is found primarily in lymphatic tissues, lymph nodes, the spleen, B lymphocytes, and macrophages (Kirschke, H. Chapter 211. Cathepsin S. In Handbook of Proteolytic Enzymes. Barrett, A. J.; Rawlings, N. D.; Woessner, J. F., Eds. San Diego: Academic Press, 1998. pp. 621-624.). Cathepsin S inhibitors have been shown in animal models to modulate antigen presentation and are effective in an animal model of asthma (Riese, R. J.; Mitchell, R. N.; Villadangos, J. A.; Shi, G.-P.; Palmer, J. T.; Karp, E. R.; De Sanctis, G. T.; Ploegh, H. L.; Chapman, H. A. Cathepsin S Activity
20 Regulates Antigen Presentation and Immunity. *J. Clin. Invest.* **1998**, *101*, 2351-2363 and Shi, G.-P.; Villadangos, J. A.; Dranoff, G.; Small, C.; Gu, L.; Haley, K. J.; Riese, R.; Ploegh, H. L.; Chapman, H. A. Cathepsin S Required for Normal MHC Class II Peptide Loading and Germinal Center Development. *Immunity* **1999**, *10*, 197-206.).

30

Mice in which the gene encoding cathepsin S has been knocked out are less susceptible to collagen-induced arthritis and their immune systems have

an impaired ability to respond to antigens (Nakagawa, T. Y.; Brissette, W. H.; Lira, P. D.; Griffiths, R. J.; Petrushova, N.; Stock, J.; McNeish, J. D.; Eastman, S. E.; Howard, E. D.; Clarke, S. R. M.; Rosloniec, E. F.; Elliott, E. A.; Rudensky, A. Y. Impaired Invariant Chain Degradation and Antigen
5 Presentation and Diminished Collagen-Induced Arthritis in Cathepsin S Null Mice. *Immunity* 1999, 10, 207-217).

These data demonstrate that compounds that inhibit the proteolytic activity of human cathepsin S should find utility in the treatment of chronic
10 autoimmune diseases including, but not limited to, lupus, rheumatoid arthritis, and asthma; and have potential utility in modulating the immune response to tissue transplantation.

There are a number of cathepsin S inhibitors reported in the literature.
15 The most important patents are listed below.

Certain dipeptidyl nitriles are claimed by Novartis as cathepsin S inhibitors in: Altmann, et. al. WO-99/24460.


20 Dipeptidyl vinyl sulfones are claimed by Arris (now Axys) as cysteine protease (including cathepsin S) inhibitors in: Palmer, et. al. US-5976858.

Certain peptidyl sulfonamides are claimed by Arris/Axys as cysteine protease (including cathepsin S) inhibitors in: Palmer, et. al. US-5776718
25 (assigned to Arris, now Axys) & Klaus, et. al. US-6030946 (assigned to Axys).

Compounds somewhat similar to those of the present invention are described in the following references.


30 Winters, et. al. (Winters, G.; Sala, A.; Barone, D.; Baldoli, E. *J. Med. Chem.* 1985, 28, 934-940; Singh, P.; Sharma, R. C. *Quant. Struct.-Act. Relat.* 1990, 9, 29-32; Winters, G.; Sala, A.; Barone, D. in US-4500525 (1985)) have described bicyclic pyrazoles of the type shown below. R never contains a

heterocyclic ring and no protease inhibitor activity is ascribed to these molecules; they are described as α 1-adrenergic receptor modulators.



5

Shutske, et. al. claim the bicyclic pyrazoles below. The pyridine ring is aromatic in their system (Shutske, G. M.; Kapples, K. J.; Tomer, J. D. US-5264576 (1993)). Although reference is made to R being a linker to a heterocycle, the claims specify only R = hydrogen. The compounds are referred to as serotonin reuptake inhibitors.




15

The compound 2-[4-[4-(3-methyl-5-phenyl-1H-pyrazol-1-yl)butyl]-1-piperazinyl]-pyrimidine is known from EP-382637, which describes pyrimidines having anxiolytic properties. This compound and analogs are further described in EP-502786 as cardiovascular and central nervous system agents. Pharmaceutical formulations with such compounds are disclosed in EP-655248 for use in the treatment of gastric secretion and as anti-ulcer agents. WO-9721439 describes medicaments with such compounds for treating obsessive-compulsive disorders, sleep apnea, sexual dysfunctions, emesis and motion sickness.

The compounds 5-methyl-3-phenyl-1-[4-(4-phenyl-1-piperazinyl)butyl]-1H-indazole and 5-bromo-3-(2-chlorophenyl)-1-[4-(4-phenyl-1-piperazinyl)butyl]-1H-indazole, in particular the hydrochloride salts thereof, are known from WO-9853940 and CA 122:314528, where these and similar
5 compounds are described as kinase inhibitors in the former reference and possessing affinity for benzodiazepine receptors in the latter reference.

Summary of the Invention

The present invention concerns compounds which can be represented by
 5 formula (I):



(I)

wherein:

the dashed line adjacent C-R⁶ is absent or an sp² bond;

Y is nitrogen or R²⁰C;

Z is nitrogen or R²¹C;

15 T is nitrogen or R²C;

S is nitrogen or R³C;

provided between 0 and 3 of S, T, Y, and Z are nitrogen; and further
 provided that one of S, T, Y, and Z can be =N⁺-O⁻ where the remaining
 three are not nitrogen;

20 R²⁰ is selected from hydrogen, halogen, C₁₋₅ alkoxy, hydroxy, C₁₋₅ alkyl,
 cyano, nitro, C₁₋₅ haloalkyl, R^oR^pN, R^oR^pNC=O, C₂₋₈ acyl, 4-7 membered
 heterocyclyl, (4-7 membered heterocyclyl)-C₁₋₅ alkylene, phenyl,
 (phenyl)C₁₋₅ alkylene, R¹⁴OC=O, R¹⁴S, R¹⁴SO, and R¹⁴SO₂;

25 R²¹ is selected from hydrogen, halogen, C₁₋₅ alkoxy, hydroxy, C₁₋₅ alkyl,
 cyano, nitro, C₁₋₅ haloalkyl, R^cR^dN, R^cR^dNC=O, C₂₋₈ acyl, 4-7 membered

heterocyclyl, (4-7 membered heterocyclyl)-C₁₋₅ alkylene, phenyl,
(phenyl)C₁₋₅ alkylene, R¹⁵OC=O, R¹⁵S, R¹⁵SO, and R¹⁵SO₂;

R² is selected from hydrogen, halogen, C₁₋₅ alkoxy, hydroxy, C₁₋₅ alkyl,
cyano, nitro, C₁₋₅ haloalkyl, R^eR^fN, R^eR^fNC=O, C₂₋₈ acyl, 4-7 membered
5 heterocyclyl, (4-7 membered heterocyclyl)-C₁₋₅ alkylene, phenyl,
(phenyl)C₁₋₅ alkylene, R¹⁶OC=O, R¹⁶S, R¹⁶SO, and R¹⁶SO₂;

R³ is selected from hydrogen, halogen, C₁₋₅ alkoxy, hydroxy, C₁₋₅ alkyl,
cyano, nitro, C₁₋₅ haloalkyl, R^gR^hN, C₂₋₈ acyl, 4-7 membered heterocyclyl,
(4-7 membered heterocyclyl)-C₁₋₅ alkylene, phenyl, (phenyl)C₁₋₅
10 alkylene, R¹⁷OC=O, R^mRⁿNC=O, R^mRⁿNSO₂, R¹⁷S, R¹⁷SO, and R¹⁷SO₂;

R⁵ and R⁶ are independently selected from hydrogen and C₁₋₅ alkyl;

R⁷ and R⁸ independently are hydrogen, C₁₋₅ alkyl, C₁₋₅ alkenyl, C₁₋₅ alkoxy, C₁₋₅
alkylthio, halogen, or 4-7 membered carbocyclyl or heterocyclyl;
alternatively, R⁷ and R⁸ can be taken together to form an optionally
15 substituted 5- to 7- membered carbocyclic or heterocyclic ring, which
ring may be unsaturated or aromatic; said ring being optionally
substituted with between 1 and 3 substituents independently selected
from halo, hydroxy, cyano, nitro, amino, R^t, R^tO-, R^tS-, R^tO(C₁₋₅
alkylene)-, R^tO(C=O)-, R^t(C=O)-, R^t(C=S)-, R^t(C=O)O-, R^tO(C=O)(C=O)-,
20 R^tSO₂, NHR^u(C=NH)-, NHR^uSO₂-, and NHR^u(C=O)-;

R^t is C₁₋₆ alkyl, phenyl, benzyl, phenethyl, or C₂₋₅ heterocyclyl, (C₁₋₅
heterocyclyl)C₁₋₆ alkylene, NH₂, mono- or di(C₁₋₆ alkyl)N-, or R⁴⁹OR⁵⁰-,
wherein R⁴⁹ is H, C₁₋₅ alkyl, C₂₋₅ alkenyl, phenyl, benzyl, phenethyl, C₁₋₅
heterocyclyl, or (C₁₋₅ heterocyclyl)C₁₋₆ alkylene and R⁵⁰ is C₁₋₅ alkylene,
25 phenylene, or divalent C₁₋₅ heterocyclyl; and

R^u can be H in addition to the values for R^t;

R^c is hydrogen, C₁₋₅ alkyl, phenyl, C₂₋₅ heterocyclyl, C₂₋₈ acyl, aroyl,
R¹⁰OC=O-, RⁱR^jNC=O, R¹⁰SO-, R¹⁰SO₂-, and RⁱR^jNSO₂;

R^e is hydrogen, C₁₋₅ alkyl, phenyl, C₂₋₅ heterocyclyl, C₂₋₈ acyl, aroyl,
30 R⁴⁰OC=O, R⁴³R⁴⁴NC=O, R⁴⁰SO, R⁴⁰SO₂, and R⁴³R⁴⁴NSO₂;

R^m is hydrogen, C₁₋₅ alkyl, phenyl, C₂₋₅ heterocyclyl, C₂₋₈ acyl, aroyl,
R⁴¹OC=O, R⁴⁵R⁴⁶NC=O, R⁴¹SO, R⁴¹SO₂, and R⁴⁵R⁴⁶NSO₂;

R^o is hydrogen, C_{1-5} alkyl, phenyl, C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl, $R^{42}OC=O$, $R^{47}R^{48}NC=O$, $R^{42}SO$, $R^{42}SO_2$, and $R^{47}R^{48}NSO_2$;

each of R^d , R^f , R^n , and R^p is independently selected from hydrogen, C_{1-5} alkyl, phenyl, and C_{2-5} heterocyclyl; in addition, R^c and R^d , R^e and R^f , R^m and R^n , or R^o and R^p , independently, can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

each of R^9 , R^{10} , R^{11} , R^{14} , R^{15} , R^{16} , R^{17} , R^{40} , R^{41} , and R^{42} is independently C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl;

each of R^i and R^j , R^k and R^l , R^{43} and R^{44} , R^{45} and R^{46} , R^{47} and R^{48} are independently hydrogen, C_{1-5} alkyl, C_{3-5} alkenyl, phenyl, or C_{2-5} heterocyclyl; in addition, R^i and R^j , and R^k and R^l , R^{43} and R^{44} , R^{45} and R^{46} , and R^{47} and R^{48} , independently, can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

R^g is hydrogen, C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl, $R^gOC=O$, $R^{18}R^{19}NC=O$, R^gSO , R^gSO_2 , or $R^{18}R^{19}NSO_2$;

R^h is hydrogen, C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl; alternatively, R^g and R^h can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

R^{18} and R^{19} independently are hydrogen, C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl; alternatively, R^{18} and R^{19} can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

n is 0, 1 or 2;

G is C_{3-6} alkenediyl or C_{3-6} alkanediyl, optionally substituted with hydroxy, halogen, C_{1-5} alkyl, C_{1-5} alkoxy, oxo, hydroximino, CO_2R^k , NR^kR^l , $(L)-C_{1-4}$ alkylene-, $R^kR^lNCO_2$, $[(L)-C_{1-5}$ alkylene]amino, N_3 , or $(L)-C_{1-5}$ alkoxy;

L is amino, mono- or di- C_{1-5} alkylamino, pyrrolidinyl, morpholinyl, piperidinyl, homopiperidinyl, or piperazinyl, wherein available ring nitrogens can be optionally substituted with C_{1-5} alkyl, benzyl, C_{2-5} acyl, C_{1-5} alkylsulfonyl, or C_{1-5} alkoxycarbonyl;

- Ar represents a monocyclic or bicyclic aryl or heteroaryl ring, optionally substituted with between 1 and 3 substituents independently selected from halogen, C₁₋₅ alkoxy, C₁₋₅ alkyl, C₂₋₅ alkenyl, cyano, azido, nitro, R²²R²³N, R²²S, R²²SO, R²²SO₂, R²²OC=O, R²²R²³NC=O, C₁₋₅ haloalkyl, C₁₋₅ haloalkoxy, C₁₋₅ haloalkylthio, and C₁₋₅ alkylthio;
- R²² is hydrogen, C₁₋₅ alkyl, C₃₋₅ alkenyl, phenyl, benzyl, C₂₋₅ heterocyclyl, C₂₋₈ acyl, aroyl, R¹¹OC=O, R²⁴R²⁵NC=O, R¹¹S, R¹¹SO, R¹¹SO₂, or R²⁴R²⁵NSO₂;
- R²³ is hydrogen, C₁₋₅ alkyl, phenyl, benzyl, or C₂₋₅ heterocyclyl;
- alternatively, R²² and R²³ can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;
- R²⁴ and R²⁵ are independently hydrogen, C₁₋₅ alkyl, phenyl, benzyl, or C₁₋₅ heteroaryl;
- alternatively, R²⁴ and R²⁵ can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;
- R³² is hydrogen, C₁₋₅ alkyl, cyano, C₁₋₅ hydroxyalkyl, C₂₋₈ acyl, -(C=O)NR^vR^x, CHO, or C₁₋₆ alkoxy carbonyl, wherein each of R^v and R^x is independently selected from H, C₁₋₅ alkyl, C₁₋₅ hydroxyalkyl, C₁₋₅ heterocyclyl, (C₁₋₅ heterocyclyl) C₁₋₅ alkylene, C₁₋₅ aminoalkylene, C₃₋₈ acyloxy, CHO, C₁₋₆ alkoxy carbonyl, and cyano;
- Q is NR³³, S, or O;
- R³³ represents hydrogen, C₁₋₅ alkyl, phenyl, benzyl, phenethyl, C₂₋₅ heterocyclyl, (C₂₋₅ heterocyclyl)C₁₋₅ alkylene, C₂₋₈ acyl, aroyl, R³⁵OC=O, R³⁶R³⁷NC=O, R³⁵SO, R³⁵S, R³⁵SO₂ and R³⁶R³⁷NSO₂;
- R³⁵ is selected from hydrogen, C₁₋₅ alkyl, phenyl, benzyl, phenethyl, and C₂₋₅ heteroaryl;
- R³⁶ and R³⁷ are each independently selected from hydrogen, C₁₋₅ alkyl, phenyl, or C₂₋₅ heteroaryl;
- alternatively, R³⁶ and R³⁷ can be taken together to form an optionally substituted 4- to 7-membered ring heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

wherein each of the above hydrocarbyl or heterocarbyl groups, unless otherwise indicated, and in addition to any specified substituents, is optionally and independently substituted with between 1 and 3 substituents selected from methyl, halomethyl, hydroxymethyl, halo, hydroxy, amino, nitro, cyano, C₁₋₅ alkyl, C₁₋₅ alkoxy, -COOH, C₂₋₆ acyl, [di(C₁₋₄ alkyl)amino]C₂₋₅ alkylene, [di(C₁₋₄ alkyl)amino] C₂₋₅ alkyl-NH-CO-, and C₁₋₅ haloalkoxy;

or a pharmaceutically acceptable salt, amide, or ester thereof; or a stereoisomeric form thereof.

The disclosed compounds are high-affinity inhibitors of the proteolytic activity of human cathepsin S. For use in medicine, the preparation of pharmaceutically acceptable salts of compounds of formula (I) may be desirable.

Certain compounds of the present invention may have one stereogenic atom and may exist as two enantiomers. Certain compounds of the present invention may have two or more stereogenic atoms and may further exist as diastereomers. It is to be understood by those skilled in the art that all such stereoisomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

Another aspect of the invention provides pharmaceutical compositions comprising a compound of formula (I) and a pharmaceutically acceptable carrier. A further embodiment of the invention is a process for making a pharmaceutical composition comprising mixing a disclosed compound as described above, with a suitable pharmaceutically acceptable carrier.

The invention also contemplates pharmaceutical compositions comprising more than one compound of formula (I) and compositions

comprising a compound of formula (I) and another pharmaceutically active agent.

5 The invention features a method of treating disorders or conditions mediated by the cathepsin S enzyme, in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above. If more than one active agent is administered, the therapeutically effective amount may be a jointly effective amount. The compounds described herein inhibit the protease
10 activity of human cathepsin S, an enzyme involved in the immune response. In preferred embodiments, cathepsin S inhibition is selective. As such, the disclosed compounds and compositions are useful in the prevention, inhibition, or treatment of autoimmune diseases such as lupus, rheumatoid arthritis, and asthma, and for the prevention, inhibition, or treatment of tissue transplant
15 rejection.

Additional features and advantages of the invention will become apparent from the detailed description below, including examples, and the appended claims.

Detailed Description of the Invention

The invention features pyrazole compounds of formula (I), methods of making them, compositions containing them, and methods of using them to treat diseases and conditions, including those mediated by Cathepsin S.

A. Terms

The following terms are defined below and by their usage throughout this disclosure.

"Alkyl" includes optionally substituted straight chain and branched hydrocarbons with at least one hydrogen removed to form a radical group. Alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, 1-methylpropyl, pentyl, isopentyl, sec-pentyl, hexyl, heptyl, octyl, and so on. Alkyl includes cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

"Alkenyl" includes optionally substituted straight chain and branched hydrocarbon radicals as above with at least one carbon-carbon double bond (sp^2). Alkenyls include ethenyl (or vinyl), prop-1-enyl, prop-2-enyl (or allyl), isopropenyl (or 1-methylvinyl), but-1-enyl, but-2-enyl, butadienyls, pentenyls, hexa-2,4-dienyl, and so on. Hydrocarbon radicals having a mixture of double bonds and triple bonds, such as 2-penten-4-ynyl, are grouped as alkynyls herein. Alkenyl includes cycloalkenyl. *Cis* and *trans* or (E) and (Z) forms are included within the invention.

"Alkynyl" includes optionally substituted straight chain and branched hydrocarbon radicals as above with at least one carbon-carbon triple bond (sp). Alkynyls include ethynyl, propynyls, butynyls, and pentynyls. Hydrocarbon radicals having a mixture of double bonds and triple bonds, such as 2-penten-4-ynyl, are grouped as alkynyls herein. Alkynyl does not include cycloalkynyl.

“Alkoxy” includes an optionally substituted straight chain or branched alkyl group with a terminal oxygen linking the alkyl group to the rest of the molecule. Alkoxy includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, pentoxy and so on. “Aminoalkyl”, “thioalkyl”, and “sulfonylalkyl” are analogous to alkoxy, replacing the terminal oxygen atom of alkoxy with, respectively, NH (or NR), S, and SO₂. Heteroalkyl includes alkoxy, aminoalkyl, thioalkyl, and so on.

“Aryl” includes phenyl, naphthyl, biphenyl, tetrahydronaphthyl, and so on, any of which may be optionally substituted. Aryl also includes arylalkyl groups such as benzyl, phenethyl, and phenylpropyl. Aryl includes a ring system containing an optionally substituted 6-membered carbocyclic aromatic ring, said system may be bicyclic, bridge, and/or fused. The system may include rings that are aromatic, or partially or completely saturated. Examples of ring systems include indenyl, pentalenyl, 1-4-dihydronaphthyl, indanyl, benzimidazolyl, benzothiophenyl, indolyl, benzofuranyl, isoquinolyl, and so on.

“Heterocyclyl” includes optionally substituted aromatic and nonaromatic rings having carbon atoms and at least one heteroatom (O, S, N) or heteroatom moiety (SO₂, CO, CONH, COO) in the ring. Unless otherwise indicated, a heterocyclic radical may have a valence connecting it to the rest of the molecule through a carbon atom, such as 3-furyl or 2-imidazolyl, or through a heteroatom, such as N-piperidyl or 1-pyrazolyl. Preferably a monocyclic heterocyclyl has between 4 and 7 ring atoms, or between 5 and 6 ring atoms; there may be between 1 and 5 heteroatoms or heteroatom moieties in the ring, and preferably between 1 and 3. A heterocyclyl may be saturated, unsaturated, aromatic (e.g., heteroaryl), nonaromatic, or fused.

30

Heterocyclyl also includes fused, e.g., bicyclic, rings, such as those optionally condensed with an optionally substituted carbocyclic or heterocyclic five- or six-membered aromatic ring. For example, “heteroaryl” includes an

optionally substituted six-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms condensed with an optionally substituted five- or six-membered carbocyclic or heterocyclic aromatic ring. Said heterocyclic five- or six-membered aromatic ring condensed with the said five- or six-membered aromatic ring may contain 1, 2 or 3 nitrogen atoms where it is a six-membered ring, or 1, 2 or 3 heteroatoms selected from oxygen, nitrogen and sulfur where it is a five-membered ring.

Examples of heterocyclyls include thiazoylyl, furyl, pyranlyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl. For example, preferred heterocyclyls or heterocyclic radicals include morpholinyl, piperazinyl, pyrrolidinyl, pyridyl, cyclohexylimino, cycloheptylimino, and more preferably, piperidyl.

Examples illustrating heteroaryl are thienyl, furanyl, pyrrolyl, imidazolyl, oxazolyl, thiazolyl, benzothienyl, benzofuranyl, benzimidazolyl, benzoxazolyl, benzothiazolyl.

"Acyl" refers to a carbonyl moiety attached to either a hydrogen atom (i.e., a formyl group) or to an optionally substituted alkyl or alkenyl chain, or heterocyclyl.

"Halo" or "halogen" includes fluoro, chloro, bromo, and iodo, and preferably chloro or bromo as a substituent.

"Alkanediyl" or "alkylene" represents straight or branched chain optionally substituted bivalent alkane radicals such as, for example, methylene, ethylene, propylene, butylene, pentylene or hexylene.

"Alkenediyl" represents, analogous to the above, straight or branched chain optionally substituted bivalent alkene radicals such as, for example, propenylene, butenylene, pentenylene or hexenylene. In such radicals, the carbon atom linking a nitrogen preferably should not be unsaturated.

5 "Aroyl" refers to a carbonyl moiety attached to an optionally substituted aryl or heteroaryl group, wherein aryl and heteroaryl have the definitions provided above. In particular, benzoyl is phenylcarbonyl.

10 As defined herein, two radicals, together with the atom(s) to which they are attached may form an optionally substituted 4- to 7-, 5- to 7-, or a 5- to 6-membered ring carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic. Said rings may be as defined above in the Summary of the Invention section. Particular examples of such rings are as follows in the next section.

15

"Pharmaceutically acceptable salts, esters, and amides" include carboxylate salts (e.g., C₁₋₆ alkyl, cycloalkyl, aryl, heteroaryl, or non-aromatic heterocyclic) amino acid addition salts, esters, and amides which are within a reasonable benefit/risk ratio, pharmacologically effective and suitable for
20 contact with the tissues of patients without undue toxicity, irritation, or allergic response. Representative salts include hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and
25 laurylsulfonate. These may include alkali metal and alkali earth cations such as sodium, potassium, calcium, and magnesium, as well as non-toxic ammonium; quaternary ammonium, and amine cations such as tetramethyl ammonium, methylamine, trimethylamine, and ethylamine. See example, S.M. Berge, et al., "Pharmaceutical Salts," J. Pharm. Sci., 1977, 66:1-19 which is
30 incorporated herein by reference. Representative pharmaceutically acceptable amides of the invention include those derived from ammonia, primary C₁₋₆ alkyl amines and secondary di (C₁₋₆ alkyl) amines. Secondary amines include 5- or 6-membered heterocyclic or heteroaromatic ring moieties containing at least

one nitrogen atom and optionally between 1 and 2 additional heteroatoms. Preferred amides are derived from ammonia, C₁₋₃ alkyl primary amines, and di (C₁₋₂ alkyl)amines. Representative pharmaceutically acceptable esters of the invention include C₁₋₇ alkyl, C₅₋₇ cycloalkyl, phenyl, and phenyl(C₁₋₆)alkyl
5 esters. Preferred esters include methyl esters.

"Patient" or "subject" includes mammals such as humans and animals (dogs, cats, horses, rats, rabbits, mice, non-human primates) in need of observation, experiment, treatment or prevention in connection with the
10 relevant disease or condition. Preferably, the patient or subject is a human.

"Composition" includes a product comprising the specified ingredients in the specified amounts as well as any product which results directly or indirectly from combinations of the specified ingredients in the specified
15 amounts.

"Therapeutically effective amount" or "effective amount" means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by
20 a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

Concerning the various radicals in this disclosure and in the claims, three general remarks are made. The first remark concerns valency. As with
25 all hydrocarbon radicals, whether saturated, unsaturated or aromatic, and whether or not cyclic, straight chain, or branched, and also similarly with all heterocyclic radicals, each radical includes substituted radicals of that type and monovalent, bivalent, and multivalent radicals as indicated by the context of the claims. The context will indicate that the substituent is an alkylene or
30 hydrocarbon radical with at least two hydrogen atoms removed (bivalent) or more hydrogen atoms removed (multivalent). An example of a bivalent radical linking two parts of the molecule is G in formula (I) which links two rings.

Second, radicals or structure fragments as defined herein are understood to include substituted radicals or structure fragments. Hydrocarbyls include monovalent radicals containing carbon and hydrogen such as alkyl, alkenyl, alkynyl, cycloalkyl, and cycloalkenyl (whether aromatic or unsaturated), as well as corresponding divalent radicals such as alkylene, alkenylene, phenylene, and so on. Heterocarbyls include monovalent and divalent radicals containing carbon, hydrogen, and at least one heteroatom. Examples of monovalent heterocarbyls include acyl, acyloxy, alkoxyacyl, heterocyclyl, heteroaryl, aroyl, benzoyl, dialkylamino, hydroxyalkyl, and so on. Using "alkyl" as an example, "alkyl" should be understood to include substituted alkyl having one or more substitutions, such as between 1 and 5, 1 and 3, or 2 and 4 substituents. The substituents may be the same (dihydroxy, dimethyl), similar (chlorofluoro), or different (chlorobenzyl- or aminomethyl-substituted). Examples of substituted alkyl include haloalkyl (such as fluoromethyl, chloromethyl, difluoromethyl, perchloromethyl, 2-bromoethyl, perfluoromethyl, and 3-iodocyclopentyl), hydroxyalkyl (such as hydroxymethyl, hydroxyethyl, 2-hydroxypropyl, aminoalkyl (such as aminomethyl, 2-aminoethyl, 3-aminopropyl, and 2-aminopropyl), nitroalkyl, alkylalkyl, and so on. A di(C₁₋₆ alkyl)amino group includes independently selected alkyl groups, to form, for example, methylpropylamino and isopropylmethylamino, in addition dialkylamino groups having two of the same alkyl group such as dimethyl amino or diethylamino.

Third, only stable compounds are intended. For example, where there is an NR'R" group, and R can be an alkenyl group, the double bond is at least one carbon removed from the nitrogen to avoid enamine formation. Similarly, where a dashed line is an optional sp² bond, if it is absent, the appropriate hydrogen atom(s) is(are) included.

Preferred substitutions for Ar include methyl, methoxy, fluoromethyl, difluoromethyl, perfluoromethyl (trifluoromethyl), 1-fluoroethyl, 2-fluoroethyl, ethoxy, fluoro, chloro, and bromo, and particularly methyl; bromo, chloro, perfluoromethyl, perfluoromethoxy, methoxy, and fluoro. Preferred substitution
5 patterns for Ar are 4-substituted or 3,4-disubstituted phenyl. Compounds of the invention are further described in the next section.

B. Compounds

The invention features compounds of formula (I) as described in the
5 Summary section.

Preferred compounds include those wherein:

- (a) one of S, T, Y, and Z is nitrogen;
- 10 (b) S and T are CR³ and CR², respectively;
- (c) S, T, Y, and Z are CR³, CR², CR²⁰, and CR²¹, respectively;
- (d) (1) Z is N, Y is N, S is CR³, and T is CR²; or (2) S is N, T is N, Y is
CR²⁰, and Z is CR²¹;
- (e) R² is hydrogen, halogen, C₁₋₅ alkoxy, cyano, R^eR^fN, or a 5-6
15 membered heterocyclyl;
- (f) R³ is hydrogen, halogen, C₁₋₅ alkoxy, C₁₋₅ alkyl, cyano, R¹⁷OC=O, or
R^gR^hN, where R^g and R^h are H or C₁₋₅ alkyl, or are taken together to
form a 5-6 membered heterocyclyl;
- (g) each of R² and R³ is independently selected from hydrogen, halogen,
20 and a 5-6 membered heterocyclyl;
- (h) R⁵ and R⁶ are independently selected from hydrogen and C₁₋₃ alkyl;
- (i) one of R⁵ and R⁶ is H;
- (j) R⁵ and R⁶ are each H;
- (k) one of R⁷ and R⁸ is H and the other is 5-7 membered carbocyclyl or
25 heterocyclyl;
- (l) R⁷ and R⁸ are taken together to form an optionally substituted 5- to
7- membered carbocyclic or heterocyclic ring;
- (m) R⁷ and R⁸ are taken together to form a six-membered heterocyclyl;
- (n) R⁷ and R⁸ taken together form a 5-7 membered heterocyclyl
30 optionally N-substituted with R^t(C=O)-, R^tSO₂-, or NHR^u(C=O)-
wherein R^t is C₁₋₆ alkyl, phenyl, or C₂₋₅ heterocyclyl and R^u is H, C₁₋₆
alkyl, phenyl, or C₂₋₅ heterocyclyl;

- (o) each of R^c , R^e , R^m , and R^o is independently selected from hydrogen, C_{1-5} alkyl, C_{2-8} acyl, $(C_{1-5} \text{ alkyl})OC=O$, and the respective $RRNC=O$, RSO , RSO_2 , and $RRNSO_2$ groups;
- (p) each of R^c , R^d , R^g , R^h , R^o , R^f , and R^p is independently selected from hydrogen and C_{1-5} alkyl; or, independently, R^e and R^f , R^g and R^h , or R^o and R^p taken together form an optionally substituted 4- to 7-membered carbocyclic or heterocyclic ring;
- (q) R^e and R^f taken together are morpholinyl, piperidinyl, or pyrrolidinyl;
- (r) each of R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^i , R^j , R^k and R^l independently is hydrogen or C_{1-5} alkyl;
- (s) each of R^9 , R^{11} , R^{14} , R^{15} , R^{16} and R^{17} is independently C_{1-5} alkyl;
- (t) R^9 is C_{1-5} alkyl, C_{2-8} acyl, $R^9OC=O$, $R^{18}R^{19}NC=O$, R^9SO , R^9SO_2 , or $R^{18}R^{19}NSO_2$; and R^h is H or C_{1-5} alkyl; alternatively, R^9 and R^h can be taken together to form an optionally substituted 5- to 6-membered heterocyclyl;
- (u) R^9 and R^h are each C_{1-3} alkyl;
- (v) R^{18} and R^{19} independently are hydrogen or C_{1-5} alkyl;
- (w) n is 0 or 1; or n is 1;
- (x) G is C_{3-4} alkanediyl, optionally substituted with hydroxy, halogen, $[(L)-C_{1-5} \text{ alkylene}]amino$, or $(L)-C_{1-5} \text{ alkyloxy}$;
- (y) G is C_3 alkanediyl, optionally substituted with hydroxy;
- (z) R^{20} and R^{21} are independently selected from hydrogen, halogen, C_{1-5} alkoxy, C_{1-5} alkyl, cyano, nitro, 4-7 membered heterocyclyl, and R^oR^pN or R^cR^dN , respectively;
- (aa) R^{20} and R^{21} are independently selected from hydrogen, halogen, 5- to 6-membered heterocyclyl, and R^oR^pN or R^cR^dN , respectively;
- (bb) Ar represents a monocyclic ring, optionally substituted with 1 to 2 substituents selected from halogen, C_{1-5} alkyl, cyano, nitro, $R^{22}R^{23}N$, C_{1-3} haloalkyl, and C_{1-3} haloalkoxy;
- (cc) Ar is a six-membered aromatic ring monosubstituted at the 4-position with halogen, methyl, CF_3 , or OCF_3 , or disubstituted at the 3- and 4-positions with substituents independently selected from halogen, CF_3 , methyl, and OCF_3 ;

(dd) each of R^{22} , R^{23} , and R^{24} is independently hydrogen or C_{1-5} alkyl;

(ee) R^{25} and R^{26} independently are hydrogen or C_{1-5} alkyl,

or, alternatively, R^{25} and R^{26} are taken together to form an optionally substituted 4- to 7- membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

(ff) R^{25} and R^{26} independently are hydrogen or C_{1-5} alkyl;

(gg) Q is NR^{33} or S;

(hh) Q is NR^{33} , R^{33} is H or C_{2-5} heterocyclyl, and R^{32} is H, C_{1-5} alkyl, C_{1-5} hydroxyalkyl, $-(C=O)NR^vR^x$, CHO, or C_{1-6} alkoxy carbonyl, wherein each of R^v and R^x is independently selected from H, C_{1-5} hydroxyalkyl, $(C_{1-5}$ heterocyclyl)- C_{1-5} alkylene, and C_{1-5} aminoalkylene;

(ii) wherein Q is S and R^{33} is $NR^{36}R^{37}(C=O)-$ where each of R^{36} and R^{37} are independently selected from hydrogen and C_{1-5} alkyl;

(jj) R^{35} is selected from hydrogen and C_{1-5} alkyl; R^{36} and R^{37} are each independently selected from hydrogen, C_{1-5} alkyl, or, alternatively, R^{36} and R^{37} can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring;

(kk) Y is nitrogen or $R^{20}C$; Z is nitrogen or $R^{21}C$; T is nitrogen or $R^{22}C$;

S is nitrogen or R^3C ; provided between 0 and 2 of S, T, Y, and Z are nitrogen; for example 1 of them is N;

(ll) R^2 is hydrogen, halogen, hydroxy, C_{1-5} alkoxy, C_{1-5} alkyl, 5- to 6-membered heterocyclyl, or R^eR^fN ;

(mm) R^3 is hydrogen, halogen, C_{1-5} alkoxy, hydroxy, C_{1-5} alkyl, , 5- to 6-membered heterocyclyl, or R^gR^hN ;

(nn) R^7 and R^8 independently are taken together to form an optionally substituted 5- to 7- membered unsaturated heterocyclic ring;

(oo) each of R^a , R^e , R^m , and R^o is independently selected from hydrogen, C_{1-5} alkyl, C_{2-8} acyl, $(C_{1-5}$ alkyl)OC=O, and the respective $RRNC=O$, RSO , RSO_2 , and $RRNSO_2$ groups;

(pp) each of R^b , R^f , R^n , and R^p , is independently selected from hydrogen and C_{1-5} alkyl; each of R^9 , R^{11} , R^{14} , R^{15} , R^{16} , R^{17} , R^{40} , R^{41} and R^{42} is independently C_{1-5} alkyl; and each of R^c , R^d , R^i , R^j , R^{43} ,

R^{44} , R^{45} , R^{46} , R^{47} , R^k and R^l are independently are hydrogen or C_{1-5} alkyl;

(qq) R^g is hydrogen, or C_{1-5} alkyl, C_{2-8} acyl, $R^gOC=O$, $R^{18}R^{19}NC=O$, R^gSO , R^gSO_2 , or $R^{18}R^{19}NSO_2$; R^h is hydrogen or C_{1-5} alkyl;

5 alternatively, R^g and R^h can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic; R^{18} and R^{19} independently are hydrogen or C_{1-5} alkyl; n is 0 or 1;

10 (rr) G is C_{3-4} alkenediyl or C_{3-4} alkanediyl, optionally substituted with hydroxy, halogen, C_{1-5} alkyloxy, oxo, hydroximino, CO_2R^k , $R^kR^lNCO_2$, or (L)- C_{1-5} alkoxy; L is amino, mono- or di- C_{1-5} alkylamino, pyrrolidinyl, morpholinyl, piperidinyl homopiperidinyl, or piperazinyl, available ring nitrogens being optionally with C_{1-5} alkyl, benzyl, C_{2-5} acyl, or C_{1-5} alkyloxycarbonyl;

15 (ss) R^{20} and R^{21} are independently selected from hydrogen, halogen, C_{1-5} alkoxy, C_{1-5} alkyl, cyano, nitro, and R^oR^pN ; alternatively, R^3 and R^{20} or R^3 and R^{21} can be taken together to form an optionally substituted 5- or 6-membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic; and Ar represents a
20 monocyclic or bicyclic aryl or heteroaryl ring, optionally substituted with hydrogen, halogen, C_{1-5} alkoxy, C_{1-5} alkyl, cyano, nitro, $R^{22}R^{23}N$, $R^{24}SO_2$, $R^{24}OC=O$, $R^{25}R^{26}NC=O$, CF_3 , OCF_3 , SCF_3 , or C_{1-5} alkylthio; R^{22} is hydrogen, C_{1-5} alkyl, phenyl, benzyl, phenethyl, C_{2-5} heteroaryl, C_{2-8} acyl, aroyl, $R^{24}OC=O$, $R^{25}R^{26}NC=O$, $R^{24}SO$, $R^{24}SO_2$, or
25 $R^{25}R^{26}NSO_2$; R^{23} is hydrogen or C_{1-5} alkyl; alternatively, R^{22} and R^{23} can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic; R^{24} is hydrogen or C_{1-5} alkyl; R^{25} and R^{26} are independently hydrogen or C_{1-5} alkyl; or, alternatively, R^{25}
30 and R^{26} can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

- (tt) R^{32} is hydrogen, C_{1-5} alkyl, C_{1-5} hydroxyalkyl, CHO, C_{2-6} acyl, C_{1-6} alkoxy carbonyl, or $-(C=O)NR^yR^x$, wherein each of R^y R^x is independently selected from H, C_{1-5} alkyl, C_{1-5} hydroxyalkyl, C_{3-8} acyloxy, (amino) C_{1-6} alkylene, (C_{1-5} heterocyclyl) C_{1-5} alkylene, or C_{1-6} alkoxy carbonyl; and Q is NR^{33} or S; R^{33} represents hydrogen, C_{1-5} alkyl, phenyl, benzyl, (C_{2-5} heterocyclyl) C_{1-5} alkylene, C_{2-6} acyl, aroyl, $R^{35}OC=O$, $R^{36}R^{37}NC=O$, $R^{35}SO_2$ and $R^{36}R^{37}NSO_2$; R^{35} is selected from hydrogen and C_{1-5} alkyl; R^{36} and R^{37} are each independently selected from hydrogen and C_{1-5} alkyl;
- (uu) one of R^5 and R^6 is H, R^7 and R^8 are taken together to form an optionally substituted 6-membered carbocyclic or heterocyclic ring; and Ar represents a monocyclic ring, optionally substituted with 1 to 2 substituents selected from halogen, C_{1-5} alkyl, cyano, nitro, $R^{22}R^{23}N$, CF_3 and OCF_3 ;
- (vv) both R^5 and R^6 are each H, and
- (ww) Ar is a six membered ring substituted with halogen, CF_3 , methyl, halomethyl, or OCF_3 , at the 3- or 4- position, or disubstituted at the 3- and 4- positions;
- (xx) R^7 and R^8 taken together form pyridinyl, pyrimidinyl, or piperazinyl, optionally N-substituted with $-(C=O)R^t$, SO_2-R^t , or $-(C=O)NHR^u$;
- (yy) R^e and R^f taken together are independently morpholinyl, piperidyl, or pyrrolidinyl, optionally substituted;
- (zz) the dashed line adjacent C- R^6 is absent;
- (aaa) or combinations of the above.

Specific preferred compounds include those in the Examples below, such as:

- 1-[1-{2-Hydroxy-3-[4-(1H-indol-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone ; 1-[4-(5-Fluoro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ; 1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-

yl]-3-[4-(5-chloro-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ; 1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-chloro-2-methyl-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-methyl-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ; 3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-carbonitrile ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-methoxy-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ; 3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-carboxylic acid ethyl ester ; 1-[4-(6-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ; 1-[1-(3-{4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-2-hydroxy-propyl)-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ; 1-[4-(5-Dimethylamino-1H-pyrrolo[3,2-b]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ; 1-[4-(5-Dimethylamino-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ; 3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-pyrrolo[2,3-b]pyridine-6-carbonitrile ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl]-propan-2-ol ; 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-morpholin-4-yl-1H-

- pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ; 1-[4-(6-Fluoro-2-hydroxymethyl-benzo[b]thiophen-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ; 6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carbaldehyde ; 6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid methyl ester ; 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid amide ; and 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic.
- 15 Furthermore, preferred compounds include those wherein Ar is selected from 4-trifluoromethylphenyl, 4-bromophenyl, 4-chlorophenyl, 4-chloro-3-methylphenyl and 3,4-dichlorophenyl.

More preferred compounds include those in Examples 4, 9, 13, and 26.

Related Compounds

5 The invention provides the disclosed compounds and closely related, pharmaceutically acceptable forms of the disclosed compounds, such as salts, esters, amides, acids, hydrates or solvated forms thereof; masked or protected forms; and racemic mixtures, or enantiomerically or optically pure forms.

Related compounds also include compounds of the invention that have been modified to be detectable, e.g., isotopically labelled with ^{18}F for use as a probe
10 in positron emission tomography (PET) or single-photon emission computed tomography (SPECT).

The invention also includes disclosed compounds having one or more functional groups (e.g., hydroxyl, amino, or carboxyl) masked by a protecting
15 group. See, e.g., Greene and Wuts, Protective Groups in Organic Synthesis, 3rd ed., (1999) John Wiley & Sons, NY. Some of these masked or protected compounds are pharmaceutically acceptable; others will be useful as intermediates. Synthetic intermediates and processes disclosed herein, and minor modifications thereof, are also within the scope of the invention.

20

HYDROXYL PROTECTING GROUPS

Protection for the hydroxyl group includes methyl ethers, substituted methyl ethers, substituted ethyl ethers, substitute benzyl ethers, and silyl ethers.

Substituted Methyl Ethers

Examples of substituted methyl ethers include methoxymethyl, methylthiomethyl, *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl, benzyloxymethyl, *p*-methoxybenzyloxymethyl, (4-methoxyphenoxy)methyl, guaiacolmethyl, *t*-butoxymethyl, 4-pentenylloxymethyl, siloxymethyl, 2-methoxyethoxymethyl, 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl, tetrahydropyranyl, 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl, 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl *S,S*-dioxido, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl, 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl and 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl.

Substituted Ethyl Ethers

Examples of substituted ethyl ethers include 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, and benzyl.

Substituted Benzyl Ethers

Examples of substituted benzyl ethers include *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2- and 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p*, *p'*-dinitrobenzhydryl, 5-dibenzosuberyl,

triphenylmethyl, α -naphthylidiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxy)phenyldiphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(1-imidazol-1-ylmethyl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-2-yl, and benzisothiazolyl S,S-dioxido.

Silyl Ethers

10 Examples of silyl ethers include trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylisopropylsilyl, diethylisopropylsilyl, dimethylthexylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl, and *t*-butylmethoxyphenylsilyl.

Esters

15 In addition to ethers, a hydroxyl group may be protected as an ester. Examples of esters include formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, *p*-*P*-phenylacetate, 3-phenylpropionate, 4-oxopentanoate(levulinate), 4,4-
20 (ethylenedithio)pentanoate, pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate(mesitoate)

Carbonates

25 Examples of carbonate protecting groups include methyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 2-(trimethylsilyl)ethyl, 2-(phenylsulfonyl)ethyl, 2-(triphenylphosphonio)ethyl, isobutyl, vinyl, allyl, *p*-nitrophenyl, benzyl, *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, S-benzyl thiocarbonate, 4-ethoxy-1-naphthyl, and methyl
30 dithiocarbonate.

Assisted Cleavage

- Examples of assisted cleavage include 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl)benzoate, 2-
- 5 formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl carbonate, 4-(methylthiomethoxy)butyrate, and 2-(methylthiomethoxymethyl)benzoate.

Miscellaneous Esters

- 10 Examples of miscellaneous esters include 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (E)-2-methyl-2-
- 15 butenoate(tigloate), *o*-(methoxycarbonyl)benzoate, *p*-P-benzoate, α -naphthoate, nitrate, alkyl N,N,N',N'-tetramethylphosphorodiamidate, N-phenylcarbamate, borate, dimethylphosphinothioyl, and 2,4-dinitrophenylsulfenate.

Sulfonates

- 20 Examples of sulfonates include sulfate, methanesulfonate(mesylate), benzylsulfonate, and tosylate.

25 AMINO PROTECTING GROUPS

Protection for the amino group includes carbamates, amides, and special -NH protective groups.

- 30 Examples of carbamates include methyl and ethyl carbamates, substituted ethyl carbamates, assisted cleavage carbamates, photolytic cleavage carbamates, urea-type derivatives, and miscellaneous carbamates.

Carbamates

Examples of methyl and ethyl carbamates include methyl and ethyl, 9-fluorenylmethyl, 9-(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl, and 4-methoxyphenacyl.

Substituted Ethyl

Examples of substituted ethyl carbamates include 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-phenylethyl, 1-(1-adamantyl)-1-methylethyl, 1,1-dimethyl-2-haloethyl, 1,1-dimethyl-2,2-dibromoethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 1-methyl-1-(4-biphenyl)ethyl, 1-(3,5-di-*t*-butylphenyl)-1-methylethyl, 2-(2'- and 4'-pyridyl)ethyl, 2-(*N,N*-dicyclohexylcarboxamido)ethyl, *t*-butyl, 1-adamantyl, vinyl, allyl, 1-isopropylallyl, cinnamyl, 4-nitrocinnamyl, 8-quinolyl, *N*-hydroxypiperidinyl, alkyledithio, benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylsulfinylbenzyl, 9-anthrylmethyl and diphenylmethyl.

Assisted Cleavage

Examples of assisted cleavage include 2-methylthioethyl, 2-methylsulfonylethyl, 2-(*p*-toluenesulfonyl)ethyl, [2-(1,3-dithianyl)]methyl, 4-methylthiophenyl, 2,4-dimethylthiophenyl, 2-phosphonioethyl, 2-triphenylphosphonioisopropyl, 1,1-dimethyl-2-cyanoethyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, 5-benzisoxazolylmethyl, and 2-(trifluoromethyl)-6-chromonylmethyl.

Photolytic Cleavage

Examples of photolytic cleavage include *m*-nitrophenyl, 3,5-dimethoxybenzyl, *o*-nitrobenzyl, 3,4-dimethoxy-6-nitrobenzyl, and phenyl(*o*-nitrophenyl)methyl.

Urea-Type Derivatives

Examples of urea-type derivatives include phenothiazinyl-(10)-carbonyl derivative, N' -*p*-toluenesulfonylaminocarbonyl, and N'-
5 phenylaminothiocarbonyl.

Miscellaneous Carbamates

Examples of miscellaneous carbamates include *t*-amyl, S-benzyl
10 thiocarbamate, *p*-cyanobenzyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropylmethyl, *p*-decyloxybenzyl, diisopropylmethyl, 2,2-dimethoxycarbonylvinyl, *o*-(N,N-dimethylcarboxamido)benzyl, 1,1-dimethyl-3-(N,N-dimethylcarboxamido)propyl, 1,1-dimethylpropynyl, di(2-pyridyl)methyl, 2-furanylmethyl, 2-iodoethyl, isobornyl, isobutyl, isonicotinyl, *p*-(*p*'-
15 methoxyphenylazo)benzyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1-methyl-1-cyclopropylmethyl, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl, 1-methyl-1-(*p*-phenylazophenyl)ethyl, 1-methyl-1-phenylethyl, 1-methyl-1-(4-pyridyl)ethyl, phenyl, *p*-(phenylazo)benzyl, 2,4,6-tri-*t*-butylphenyl, 4-(trimethylammonium)benzyl, and 2,4,6-trimethylbenzyl.

20

Examples of amides include:

Amides

N-formyl, N-acetyl, N-chloroacetyl, N-trichloroacetyl, N-trifluoroacetyl, N-
25 phenylacetyl, N-3-phenylpropionyl, N-picolinoyl, N-3-pyridylcarboxamide, N-benzoylphenylalanyl derivative, N-benzoyl, N-*p*-phenylbenzoyl.

Assisted Cleavage

N-*o*-nitrophenylacetyl, N-*o*-nitrophenoxyacetyl, N-acetoacetyl, (N'-
30 dithiobenzyloxycarbonylamino)acetyl, N-3-(*p*-hydroxyphenyl)propionyl, N-3-(*o*-nitrophenyl)propionyl, N-2-methyl-2-(*o*-nitrophenoxy)propionyl, N-2-methyl-2-(*o*-phenylazophenoxy)propionyl, N-4-chlorobutyryl, N-3-methyl-3-nitrobutyryl, N-*o*-nitrocinnamoyl, N-acetylmethionine derivative, N-*o*-nitrobenzoyl, N-*o*-(benzoyloxymethyl)benzoyl, and 4,5-diphenyl-3-oxazolin-2-one.

Cyclic Imide Derivatives

5 N-phthalimide, N-dithiasuccinoyl, N-2,3-diphenylmaleoyl, N-2,5-dimethylpyrrolyl, N-1,1,4,4-tetramethyldisilylazacyclopentane adduct, 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, and 1-substituted 3,5-dinitro-4-pyridonyl.

10

SPECIAL – NH PROTECTIVE GROUPS

Examples of special NH protective groups include

15 N-Alkyl and N-Aryl Amines

N-methyl, N-allyl, N-[2-(trimethylsilyl)ethoxy]methyl, N-3-acetoxypentyl, N-(1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl), quaternary ammonium salts, N-benzyl, N-di(4-methoxyphenyl)methyl, N-5-dibenzosuberyl, N-triphenylmethyl, 20 N-(4-methoxyphenyl)diphenylmethyl, N-9-phenylfluorenyl, N-2,7-dichloro-9-fluorenylmethylene, N-ferrocenylmethyl, and N-2-picolylamine N'-oxide.

Imine Derivatives

25 N-1,1-dimethylthiomethylene, N-benzylidene, N-*p*-methoxybenzylidene, N-diphenylmethylene, N-[(2-pyridyl)mesityl]methylene, and N-(N',N'-dimethylaminomethylene).

30 PROTECTION FOR THE CARBONYL GROUP

Acyclic Acetals and Ketals

Examples of acyclic acetals and ketals include dimethyl, bis(2,2,2-trichloroethyl), dibenzyl, bis(2-nitrobenzyl) and diacetyl.

35

Cyclic Acetals and Ketals

Examples of cyclic acetals and ketals include 1,3-dioxanes, 5-methylene-1,3-dioxane, 5,5-dibromo-1,3-dioxane, 5-(2-pyridyl)-1,3-dioxane, 1,3-dioxolanes, 4-bromomethyl-1,3-dioxolane, 4-(3-butenyl)-1,3-dioxolane, 4-phenyl-1,3-dioxolane, 4-(2-nitrophenyl)-1,3-dioxolane, 4,5-dimethoxymethyl-1,3-dioxolane, O,O'-phenylenedioxy and 1,5-dihydro-3H-2,4-benzodioxepin.

Acyclic Dithio Acetals and Ketals

Examples of acyclic dithio acetals and ketals include S,S'-dimethyl, S,S'-diethyl, S,S'-dipropyl, S,S'-dibutyl, S,S'-dipentyl, S,S'-diphenyl, S,S'-dibenzyl and S,S'-diacetyl.

Cyclic Dithio Acetals and Ketals

Examples of cyclic dithio acetals and ketals include 1,3-dithiane, 1,3-dithiolane and 1,5-dihydro-3H-2,4-benzodithiepin.

Acyclic Monothio Acetals and Ketals

Examples of acyclic monothio acetals and ketals include O-trimethylsilyl-S-alkyl, O-methyl-S-alkyl or -S-phenyl and O-methyl-S-2-(methylthio)ethyl.

Cyclic Monothio Acetals and Ketals

Examples of cyclic monothio acetals and ketals include 1,3-oxathiolanes.

MISCELLANEOUS DERIVATIVES

O-Substituted Cyanohydrins

Examples of O-substituted cyanohydrins include O-acetyl, O-trimethylsilyl, O-1-ethoxyethyl and O-tetrahydropyranyl.

Substituted Hydrazones

Examples of substituted hydrazones include N,N-dimethyl and 2,4-dinitrophenyl.

5

Oxime Derivatives

Examples of oxime derivatives include O-methyl, O-benzyl and O-phenylthiomethyl.

10

Imines

Substituted Methylene Derivatives, Cyclic Derivatives

15

Examples of substituted methylene and cyclic derivatives include oxazolidines, 1-methyl-2-(1'-hydroxyalkyl)imidazoles, N,N'-dimethylimidazolidines, 2,3-dihydro-1,3-benzothiazoles, diethylamine adducts, and methylaluminum bis(2,6-di-*t*-butyl-4-methylphenoxide)(MAD)complex.

PROTECTION FOR THE CARBOXYL GROUP

Esters

5 Substituted Methyl Esters

Examples of substituted methyl esters include 9-fluorenylmethyl, methoxymethyl, methylthiomethyl, tetrahydropyranyl, tetrahydrofuranyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, benzyloxymethyl,
10 phenacyl, *p*-bromophenacyl, α -methylphenacyl, *p*-methoxyphenacyl, carboxamidomethyl, and N-phthalimidomethyl.

2-Substituted Ethyl Esters

15 Examples of 2-substituted ethyl esters include 2,2,2-trichloroethyl, 2-haloethyl, ω -chloroalkyl, 2-(trimethylsilyl)ethyl, 2-methylthioethyl, 1,3-dithianyl-2-methyl, 2-(*p*-nitrophenylsulfenyl)ethyl, 2-(*p*-toluenesulfonyl)ethyl, 2-(2'-pyridyl)ethyl, 2-(diphenylphosphino)ethyl, 1-methyl-1-phenylethyl, *t*-butyl, cyclopentyl, cyclohexyl, allyl, 3-buten-1-yl, 4-(trimethylsilyl)-2-buten-1-yl,
20 cinnamyl, α -methylcinnamyl, phenyl, *p*-(methylmercapto)phenyl and benzyl.

Substituted Benzyl Esters

Examples of substituted benzyl esters include triphenylmethyl, 25 diphenylmethyl, bis(*o*-nitrophenyl)methyl, 9-anthrylmethyl, 2-(9,10-dioxo)anthrylmethyl, 5-dibenzosuberyl, 1-pyrenylmethyl, 2-(trifluoromethyl)-6-chromylmethyl, 2,4,6-trimethylbenzyl, *p*-bromobenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-methoxybenzyl, 2,6-dimethoxybenzyl, 4-(methylsulfinyl)benzyl, 4-sulfobenzyl, piperonyl, 4-picolyl and *p*-P-benzyl.

30

Silyl Esters

Examples of silyl esters include trimethylsilyl, triethylsilyl, *t*-butyldimethylsilyl, *i*-propyldimethylsilyl, phenyldimethylsilyl and di-*t*-
35 butylmethylsilyl.

Activated Esters

Examples of activated esters include thiols.

5

Miscellaneous Derivatives

Examples of miscellaneous derivatives include oxazoles, 2-alkyl-1,3-oxazolines, 4-alkyl-5-oxo-1,3-oxazolidines, 5-alkyl-4-oxo-1,3-dioxolanes, ortho esters, phenyl group and pentaaminocobalt(III) complex.

10

Stannyl Esters

Examples of stannyl esters include triethylstannyl and tri-*n*-butylstannyl.

15

AMIDES AND HYDRAZIDES

Amides

Examples of amides include N,N-dimethyl, pyrrolidinyl, piperidinyl, 5,6-dihydrophenanthridinyl, *o*-nitroanilides, N-7-nitroindolyl, N-8-Nitro-1,2,3,4-tetrahydroquinolyl, and *p*-P-benzenesulfonamides.

20

Hydrazides


Examples of hydrazides include N-phenyl and N,N'-diisopropyl hydrazides.

25

C. Synthesis


The compounds of the present invention may be prepared by
5 conventional synthetic organic chemistry and by matrix or combinatorial
methods according to Schemes 1 to 12 below, and Examples 1 to 13. Those
of ordinary skill in the art will be able to modify and adapt the guidance
provided herein to make the disclosed compounds.

10 Scheme 1




P = H, tert-butoxycarbonyl (BOC), EtOCO, Ac, etc.

15 Scheme 2




Scheme 3




5

Scheme 4




10

Scheme 5




5

Scheme 6




Scheme 7




5

Scheme 8




10

Scheme 9




5

Scheme 10




10

Scheme 11



Scheme 12



D. Formulation and Administration

The present compounds inhibit the proteolytic activity of human cathepsin S and therefore are useful as a medicine especially in methods for treating patients suffering from disorders or conditions which are modulated or regulated by the inhibition of cathepsin S activity.

The invention features a method for treating a subject with a condition mediated by cathepsin S, said method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of the invention. The invention also provides a method for inhibiting cathepsin S activity in a subject, wherein the method comprises administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of the invention. A third method is a method for treating an autoimmune disease, or inhibiting the progression of an autoimmune disease, in a subject, said method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a disclosed compound. The autoimmune disease can be, for example, lupus, rheumatoid arthritis, or preferably, asthma. The invention also provides a method for treating or inhibiting the progression of tissue transplant rejection in a subject, the method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of the invention. The administration step can occur before, during, and/or after a tissue transplant procedure.

In view of their inhibitory effect on the proteolytic activity of human cathepsin S the compounds of the present invention may be formulated into various pharmaceutical forms for administration purposes. To prepare these pharmaceutical compositions, an effective amount of a particular compound, in base or acid addition salt form, as the active ingredient is intimately mixed with a pharmaceutically acceptable carrier.

A carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for oral administration or parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. These include water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. In view of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are generally employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause a significant deleterious effect to the skin. Such additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment. Acid addition salts of the compounds of formula I, due to their increased water solubility over the corresponding base form, are more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce

the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated
5 multiples thereof.

Pharmaceutically acceptable acid addition salts include the therapeutically active non-toxic acid addition salt forms which the disclosed compounds are able to form. The latter can conveniently be obtained by treating the base
10 form with an appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic,
15 ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, palmoic and the like acids. The term addition salt also comprises the solvates which the disclosed compounds, as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like. Conversely the salt form can be converted by treatment with alkali
20 into the free base form.

Stereoisomeric forms defines all the possible isomeric forms which the compounds of formula (I) may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all
25 possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure. More in particular, stereogenic centers may have the (R)- or (S)-configuration; substituents on bivalent cyclic saturated radicals may have either the cis- or trans-configuration. The invention encompasses stereochemically isomeric
30 forms including diastereoisomers, as well as mixtures thereof in any proportion of the disclosed compounds. The disclosed compounds may also exist in their tautomeric forms. Such forms although not explicitly indicated in the above

and following formulae are intended to be included within the scope of the present invention.


Those of skill in the treatment of disorders or conditions mediated by the cathepsin S enzyme could easily determine the effective daily amount from the test results presented hereinafter and other information. In general it is contemplated that a therapeutically effective dose would be from 0.001 mg/kg to 5 mg/kg body weight, more preferably from 0.01 mg/kg to 0.5 mg/kg body weight. It may be appropriate to administer the therapeutically effective dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 0.05 mg to 250 mg, and in particular 0.5 to 50 mg of active ingredient per unit dosage form. Examples include 2 mg, 4 mg, 7 mg, 10 mg, 15 mg, 25 mg, and 35 mg dosage forms. Compounds of the invention may also be prepared in time-release or subcutaneous or transdermal patch formulations. Disclosed compound may also be formulated as a spray or other topical or inhalable formulations.

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the patient may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated patient and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned herein are therefore only guidelines.

The next section includes detailed information relating to the preparation, characterization, and use of the disclosed compounds.

E. Examples

5 EXAMPLE 1



1-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

A. 1-Methanesulfonyl-piperidin-4-one.

Potassium carbonate (324 g, 2340 mmol) was added to a solution of 4-piperidone monohydrate hydrochloride (90 g, 586 mmol) in chloroform (300 mL) and water (300 mL). The slurry was cooled to 0 °C and treated with methylsulfonyl chloride (136 mL, 1760 mmol) by dropwise addition over a 1 h period (gas evolution was observed). The reaction mixture was allowed to shake for 72 h and was partitioned between CH₂Cl₂ (500 mL) and saturated aqueous NaHCO₃ (500 mL). The aqueous layer was extracted with CH₂Cl₂ (3 X 200 mL). The organic layer was washed with 1% KHSO₄ (250 mL), dried (Na₂SO₄), and concentrated to afford 90.5 g (87%) of a white solid. MS (electrospray): exact mass calculated for C₆H₁₁NO₃S, 177.1; *m/z* found, 178.1 [M+H]⁺. HPLC (reverse phase conditions): *t_R* = 2.19 min. ¹H NMR (400 MHz, CDCl₃): 3.60 (t, *J* = 6.5 Hz, 4H), 2.89 (s, 3H), 2.59 (t, *J* = 6.3 Hz, 4H).

B. 5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine.

p-Toluenesulfonic acid (1.34 g, 7.0 mmol) and morpholine (25.83 mL, 296 mmol) were added to a solution of 1-methanesulfonyl-piperidin-4-one (50.0 g, 282 mmol) in benzene (282 mL). The reaction mixture was heated in a flask

equipped with a condenser and a Dean-Stark trap at reflux for 15 h. The reaction mixture was cooled and concentrated in vacuo to give the enamine which was used without further purification. The enamine was dissolved in CH_2Cl_2 (200 mL) and cooled to 0°C . To this was added triethylamine (47.2 mL, 339 mmol) followed by dropwise addition of 4-trifluoromethylbenzoyl chloride (42.3 mL, 285 mmol) dissolved in CH_2Cl_2 (82 mL). The reaction mixture was allowed to warm to room temperature and stirred for 20 h. The reaction mixture was washed with 1 N aqueous HCl (250 mL) and the CH_2Cl_2 layer was separated, dried (Na_2SO_4), and concentrated. The resulting oil was taken up in ethanol (300 mL) and treated with hydrazine (44.3 mL, 1.41 mol) at 0°C . The reaction mixture was allowed to warm to room temperature and stirred for 24 h. The mixture was concentrated and the resulting solid was filtered with ethanol wash and dried in vacuo to afford 70 g (72%) of 5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine as a white solid. MS (electrospray): exact mass calculated for $\text{C}_{14}\text{H}_{14}\text{F}_3\text{N}_3\text{O}_2\text{S}$, 345.0; m/z found, 346.0 $[\text{M}+\text{H}]^+$. HPLC (reverse phase conditions): $t_R = 6.33$ min. ^1H NMR (400 MHz, CDCl_3): 7.72 (s, 4H), 4.58 (s, 2H), 3.69 (t, $J = 5.7$ Hz, 2H), 2.99 (t, $J = 5.7$ Hz, 2H), 2.92 (s, 3H).

C. 5-Methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine.

5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (10.0 g, 29.0 mmol) and epichlorohydrin (24 mL, 307 mmol) were set stirring in DMF (150 mL) containing Cs_2CO_3 (10.4 g, 31.9 mmol). After stirring at room temperature for 4 days the mixture was evaporated, brought up in EtOAc and washed with water. The organics were dried (MgSO_4) and evaporated to give a light yellow solid. Column chromatography (silica, 5% acetone/ CH_2Cl_2) gave 4.1 g (35%) of a white solid. TLC (silica, 5% acetone/ CH_2Cl_2): $R_f = 0.28$. MS (electrospray): exact mass calculated for $\text{C}_{17}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_3\text{S}$, 401.10; m/z found, 402.1 $[\text{M}+\text{H}]^+$. ^1H NMR (400 MHz, CDCl_3): 7.84 (d, $J = 8.3$ Hz, 2H), 7.79 (d, $J = 8.3$ Hz, 2H), 4.70-4.62 (m, 3H), 4.25 (d, $J = 5.4$ Hz, 1H), 3.90-3.70 (m, 2H), 3.47 (m, 1H), 3.10-2.9 (m, 6H), 2.65-2.60 (m, 1H).

D. 4-(5-Chloro-1H-indol-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

5-Chloro-1H-indole (3.2 g, 20 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (7.97 g, 40 mmol) and potassium hydroxide (4.5 g, 80 mmol) were added in MeOH (40 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (200 mL). The mixture were extracted with 10% MeOH/CH₂Cl₂ (5 x 100 mL). The organic extracts was dried over Na₂SO₄ and concentrated to form a solid. The solid was washed with MeOH (100 mL), filtered and dried to give a light yellow solid 6.3 g (94%). TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₁₈H₂₁ClN₂O₂, 332.12; m/z found, 355.0 [M⁺+Na]. ¹H NMR (CDCl₃, 400 MHz): 8.26 (br s, 1H), 7.83 (d, J = 1.76 Hz, 1H), 7.28 (d, J = 8.80 Hz, 1H), 7.19-7.14 (m, 2H), 6.09 (br s, 1H), 4.15-4.10 (m, 2H), 3.66 (t, J = 5.67 Hz, 2H), 2.56-2.49 (m, 2H), 1.50 (s, 9H).

E. 4-(5-Chloro-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

4-(5-Chloro-1H-indol-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (6.3 g, 18.9 mmol) in EtOH (125 mL) containing PtO₂ (1 g) was placed on a Parr hydrogenator at 60 psi H₂. After 18 h the mixture was filtered through celite and evaporated to give a white solid 6.0 g (94%). TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₁₈H₂₃ClN₂O₂, 334.14; m/z found, 335.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz) 8.46 (br s, 1H), 7.51 (d, J = 8.41 Hz, 1H), 7.32 (d, J = 1.57 Hz, 1H), 7.06 (dd, J = 6.46 Hz, 2.15 Hz, 1H), 6.92 (d, J = 2.35 Hz, 1H), 4.24 (d, J = 13.11 Hz, 2H), 2.98-2.84 (m, 3H), 2.00 (d, J = 12.72 Hz, 2H), 1.69-1.55 (m, 2H), 1.50 (s, 9H).

F. 5-Chloro-3-piperidin-4-yl-1H-indole.

4-(5-Chloro-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (3.4 g, 10.2 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated and the golden oil brought up in Et₂O. A solid formed and was filtered, washed with Et₂O and air dried to give 3.5 g (97%) of a white solid as a


TFA salt. MS (electrospray): exact mass calculated for $C_{12}H_{15}ClN_2$, 234.09; m/z found, 235.1 $[M^+ + H]$.

G. 1-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-

5 trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-
ol.

5-Chloro-3-piperidin-4-yl-1H-indole (350 mg, 1.00 mmol) and 5-methane-sulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (401 mg, 1.00 mmol) were set stirring in EtOH (20 mL) containing Et_3N (215 μL , 1.54 mmol) at 80 °C. After 16 h the mixture was cooled, evaporated, brought up in CH_2Cl_2 and washed with water. The organics were dried over Na_2SO_4 and concentrated. Column chromatography (silica, 0-10% MeOH/ CH_2Cl_2) provided 551 mg (88%) of a white solid. TLC (silica, 10% MeOH/ CH_2Cl_2): R_f = 0.8. MS (electrospray): exact mass calculated for $C_{30}H_{33}ClF_3N_5O_3S$, 635.19; m/z found, 636.2 $[M^+ + H]$. 1H NMR ($CDCl_3$, 400 MHz): 8.82 (br s, 1H), 7.68 (d, J = 8.41 Hz, 2H), 7.61 (d, J = 8.6 Hz, 2H), 7.54 (br s, 1H), 7.16 (d, J = 8.41 Hz, 1H), 7.03 (dd, J = 7.0 Hz, 1.6 Hz, 1H), 6.85 (br s, 1H), 4.43 (dd, J = 25.2 Hz, 14.6 Hz, 2H), 4.30-4.05 (m, 3H), 4.00-3.88 (m, 1H), 3.62-3.50 (m, 1H), 3.47-3.35 (m, 1H), 3.02-2.89 (m, 2H), 2.88-2.81 (m, 2H), 2.79 (s, 3H), 2.72-2.60 (m, 1H), 2.47-2.28 (m, 3H), 2.12-2.00 (m, 1H), 1.96-1.85 (m, 2H), 1.74-1.50 (m, 2H).

EXAMPLE 2



1-[4-(7-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

A. 4-(7-Chloro-1H-indol-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

7-Chloro-1H-indole (3.2g, 20 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (7.97 g, 40 mmol) and potassium hydroxide (4.5 g, 80 mmol) were added in MeOH (40 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (200 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 100 mL). The organic extracts were dried over Na₂SO₄ and concentrated to form a solid. The solid was washed with MeOH (100 mL), filtered and dried to give a light yellow solid 6.3 g (94%). TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₁₈H₂₁ClN₂O₂, 332.12; *m/z* found, 355.0 [M⁺+Na]. ¹H NMR (CDCl₃, 400 MHz): 8.26 (br s, 1H), 7.83 (d, *J* = 1.76 Hz, 1H), 7.28 (d, *J* = 8.80 Hz, 1H), 7.19-7.14 (m, 2H), 6.09 (br s, 1H), 4.15-4.10 (m, 2H), 3.66 (t, *J* = 5.67 Hz, 2H), 2.56-2.49 (m, 2H), 1.50 (s, 9H).

B. 4-(7-Chloro-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

4-(7-Chloro-1H-indol-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (6.3 g, 18.9 mmol) in EtOH (125 mL) containing PtO₂ (1 g) was placed on a Parr hydrogenator at 60 psi H₂. After 18 h the mixture was filtered through celite and evaporated to give a white solid 6.0 g (94%). TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₁₈H₂₃ClN₂O₂, 334.14; *m/z* found, 335.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.46 (br s, 1H), 7.51 (d, *J* = 8.41 Hz, 1H), 7.32 (d, *J* = 1.57 Hz, 1H), 7.06 (dd, *J* = 6.46 Hz, 2.15 Hz, 1H), 6.92 (d, *J* = 2.35 Hz, 1H), 4.24 (d, *J* = 13.11 Hz, 2H), 2.98-2.84 (m, 3H), 2.00 (d, *J* = 12.72 Hz, 2H), 1.69-1.55 (m, 2H), 1.50 (s, 9H).

C. 7-Chloro-3-piperidin-4-yl-1H-indole.


4-(7-Chloro-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (3.4 g, 10.2 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated and the golden oil brought up in Et₂O. A solid formed and was filtered, washed with Et₂O and air dried to give 3.5 g (97%) of a white solid.

MS (electrospray): exact mass calculated for $C_{12}H_{15}ClN_2$, 234.09; m/z found, 235.1 $[M^+ + H]$.

D. 1-[4-(7-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

7-Chloro-3-piperidin-4-yl-1H-indole (341 mg, 0.97 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (130 mg, 0.32 mmol) were set stirring in EtOH (15 mL) containing Et_3N (135 μ L, 0.97 mmol) at 80 °C. After 16 h the mixture was cooled, evaporated, brought up in CH_2Cl_2 and washed with water. The organics were dried over Na_2SO_4 and concentrated. Column chromatography (silica, 0-10% MeOH/ CH_2Cl_2) gave 120 mg (65%) of a white solid. TLC (silica, 10% MeOH/ CH_2Cl_2): R_f = 0.7. MS (electrospray): exact mass calculated for $C_{30}H_{33}ClF_3N_5O_3S$, 635.19; m/z found, 636.2 $[M^+ + H]$. 1H NMR ($CDCl_3$, 400 MHz): 8.55 (br s, 1H), 7.70 (d, J = 8.22 Hz, 2H), 7.63 (d, J = 8.4 Hz, 2H), 7.49 (d, J = 9.4 Hz, 1H), 7.14 (d, J = 7.8 Hz, 1H), 7.00 (t, J = 8.02 Hz, 1H), 6.94 (br s, 1H), 4.51 (dd, J = 12.5 Hz, 14.5 Hz, 2H), 4.25-4.11 (m, 3H), 4.07-3.95 (m, 1H), 3.73-3.61 (m, 1H), 3.61-3.50 (m, 1H), 3.11-2.98 (m, 2H), 2.88-2.85 (m, 2H), 2.83 (s, 3H), 2.82-2.72 (m, 1H), 2.55-2.38 (m, 3H), 2.24-2.10 (m, 1H), 2.05-1.90 (m, 2H), 1.82-1.61 (m, 2H).

EXAMPLE 3



1-[4-(5-Chloro-2-methyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

A. 4-(5-Chloro-2-methyl-1H-indol-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

5-Chloro-2-methyl-1H-indole (3.3 g, 20 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (7.97 g, 40 mmol) and potassium hydroxide (4.5 g, 80 mmol) were added in MeOH (40 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (200 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 100 mL). The organic extracts were dried over Na₂SO₄ and concentrated to form a solid. The solid was washed with MeOH (100 mL), filtered and dried to give a light yellow solid 6.2 g (90%). TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₁₉H₂₃ClN₂O₂, 346.14; m/z found, 347.1 [M⁺+H].

B. 4-(5-Chloro-2-methyl-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

4-(5-Chloro-2-methyl-1H-indol-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (6.2 g, 17.9 mmol) in EtOH (125 mL) containing PtO₂ (1 g) was placed on a Parr hydrogenator at 60 psi H₂. After 18 h the mixture was filtered through celite and evaporated to give a white solid 6.2 g (99%). TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₁₉H₂₅ClN₂O₂, 348.16; m/z found, 349.1 [M⁺+H].


C. 5-Chloro-2-methyl-3-piperidin-4-yl-1H-indole.

4-(5-Chloro-2-methyl-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (6.2 g, 107.8 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated and the golden oil brought up in Et₂O. A solid formed and was filtered, washed with Et₂O and air dried to give 6.2 g (95%) of a white solid as a TFA salt. MS (electrospray): exact mass calculated for C₁₄H₁₇ClN₂, 248.11; m/z found, 249.1 [M⁺+H].

D. 1-[4-(5-Chloro-2-methyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

5-Chloro-2-methyl-3-piperidin-4-yl-1H-indole (480 mg, 1.32 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (177 mg, 0.44 mmol) were set stirring in EtOH (20 mL) containing Et₃N (215 μ L, 1.54 mmol) at 80 °C. After 16 h the mixture was cooled, evaporated, brought up in CH₂Cl₂ and washed with water. The organics were dried over Na₂SO₄ and concentrated. Column chromatography (silica, 0-10% MeOH/CH₂Cl₂) gave 169 mg (62%) of a white solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.6. MS (electrospray): exact mass calculated for C₃₁H₃₅ClF₃N₅O₃S, 649.21; *m/z* found, 650.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.00 (s, 1H), 7.70 (d, *J* = 8.11 Hz, 2H), 7.64 (d, *J* = 8.41 Hz, 2H), 7.57 (d, *J* = 1.96 Hz, 1H), 7.12 (d, *J* = 8.61 Hz, 1H), 6.99 (dd, *J* = 6.85 Hz, 1.96 Hz, 1H), 4.53 (dd, *J* = 14.28 Hz, 12.91 Hz, 2H), 4.26-4.14 (m, 2H), 4.09-3.99 (m, 1H), 3.75-3.65 (m, 1H), 3.64-3.54 (m, 1H), 3.14-3.02 (m, 2H), 3.00-2.89 (m, 2H), 2.86 (s, 3H), 2.76-2.63 (m, 1H), 2.54-2.45 (m, 2H), 2.45-2.36 (m, 1H), 2.34 (s, 3H), 2.25-2.00 (m, 3H), 1.77-1.63 (m, 2H).

EXAMPLE 4



1-[4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

A. 6-Chloro-3-piperidin-4-yl-1H-indole.

6-Chloro-1H-indole (3.2g, 20 mmol), piperidin-4-one monohydrate (6.1 g, 40 mmol) and potassium hydroxide (4.5 g, 80 mmol) were added in MeOH (40 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (200 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 100 mL). The organic extracts were dried over Na₂SO₄ and concentrated. Column chromatography (silica, 20-100% MeOH/CH₂Cl₂ with 2% NH₄OH) to obtain 5.8 g (100%) of a yellow solid. The solid (5.8 g, 20 mmol) in EtOH (150 mL) containing PtO₂ (1 g) was placed on a Parr hydrogenator at 60 psi H₂. After 18 h the mixture was filtered through celite and evaporated to give an off white solid 4.6 g (97%). MS (electrospray): exact mass calculated for C₁₃H₁₅ClN₂, 234.09; *m/z* found, 235.0 [M⁺+H].

B. 4-(6-Chloro-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

To a solution of 6-chloro-3-piperidin-4-yl-1H-indole (4.6 g, 19.5 mmol) in DMF (20 mL) was added di-tert-butyl dicarbonate (4.6 g, 21.4 mmol). The reaction mixture was stirred at room temperature for 6 h. The reaction mixture was dissolved in EtOAc (400 mL), washed with water (3 x 50 mL), brine (1 x 50 mL). The organic layer was dried over Na₂SO₄ and concentrated. Column chromatography (silica, 20-50% EtOAc/hexanes) gave 4.2 g (64%) of the desired product. TLC (silica, 20% EtOAc/hexanes): R_f = 0.24. MS (electrospray): exact mass calculated for C₁₈H₂₃ClN₂O₂, 334.14; *m/z* found, 335.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz) 8.46 (br s, 1H), 7.42 (d, *J* = 8.61 Hz, 1H), 7.14 (d, *J* = 1.57 Hz, 1H), 6.96 (dd, *J* = 6.65 Hz, 1.76 Hz, 1H), 6.74 (s, 1H), 4.14 (br s, 2H), 2.89-2.70 (m, 3H), 1.90 (d, *J* = 12.13, 2H), 1.65-1.50 (m, 2H), 1.41 (s, 9H).

C. 4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidine-1-carboxylic acid tert-butyl ester.

4-(6-Chloro-1H-indol-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (2.0 g, 5.95 mmol) was dissolved in THF (30 mL). At 0 °C, potassium bis(trimethylsilyl)amide (2.37 g, 11.9 mmol) was added. The reaction mixture was stirred at room temperature for 1 h. 4-(2-Chloro-ethyl)-morpholine

hydrochloride (1.8 g, 11.9 mmol) was added and stirred at room temperature for an additional 1 h. The mixture was dissolved in EtOAc (250 mL) and washed with water (2 x 30 mL) and brine (30 mL). The organics were dried over Na₂SO₄ and concentrated. Column chromatography (silica, 0-5%

5 MeOH/CH₂Cl₂) provided 2.6 g (97%) of a white solid. TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.67. MS (electrospray): exact mass calculated for C₂₄H₃₄ClN₃O₃, 447.23; *m/z* found, 448.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 7.45 (d, *J* = 8.61 Hz, 1H), 7.27 (d, *J* = 1.57 Hz, 1H), 6.99 (dd, *J* = 6.65 Hz, 1.76 Hz, 1H), 6.84 (s, 1H), 4.18 (br s, 2H), 4.06 (t, *J* = 6.85 Hz, 2H), 3.69-3.60 (m, 4H), 2.92-2.80 (m, 2H), 2.69-2.60 (m, 3H), 2.44 (t, *J* = 4.89 Hz, 2H), 2.40 (t, *J* = 4.30 Hz, 2H), 1.94 (d, *J* = 12.13, 2H), 1.65-1.50 (m, 2H), 1.45 (s, 9H).

D. 6-Chloro-1-(2-morpholin-4-yl-ethyl)-3-piperidin-4-yl-1H-indole.


4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidine-1-carboxylic acid tert-butyl ester (2.6 g, 5.81 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 15 45 min the mixture was evaporated and the golden oil brought up in Et₂O. A solid formed and was filtered, washed with Et₂O and air dried to give 2.5 g (95%) of a white solid. MS (electrospray): exact mass calculated for C₁₉H₂₆ClN₃O, 347.18; *m/z* found, 348.2 [M⁺+H].

E. 1-{4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

6-Chloro-1-(2-morpholin-4-yl-ethyl)-3-piperidin-4-yl-1H-indole (209 mg, 0.6 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (120 mg, 0.3 mmol) were set stirring in EtOH (20 mL) containing Et₃N (84 μL, 0.6 mmol) at 80 °C. After 16 h the mixture was cooled, evaporated, brought up in CH₂Cl₂ and washed with water. The organics were dried over Na₂SO₄ and concentrated. Column chromatography (silica, 0-10% MeOH/CH₂Cl₂) provided 180 mg (85%) of a white solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.54. MS (electrospray): exact mass calculated for C₃₆H₄₄ClF₃N₆O₄S, 748.28; *m/z* found, 749.3 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 7.70 (d, *J* = 8.61 Hz, 2H), 7.64 (d, *J* = 8.261 Hz,

2H), 7.47 (d, $J = 8.80$ Hz, 1H), 7.29 (d, $J = 1.96$, 1H), 7.02 (dd, $J = 6.46$ Hz, 1.76 Hz, 1H), 6.81 (br s, 1H), 4.54 (dd, $J = 4.09$ Hz, 7.43 Hz, 2H), 4.24-4.14 (m, 2H), 4.14-4.08 (m, 2H), 4.06-3.98 (m, 1H), 3.73-3.57 (m, 5H), 3.12-3.02 (m, 2H), 2.97-2.87 (m, 2H), 2.86 (s, 3H), 2.83-2.74 (m, 1H), 2.70-2.64 (t, $J = 7.24$ Hz, 2H), 2.54-2.42 (m, 8H), 2.23-2.14 (m, 1H), 2.05-1.96 (m, 2H), 1.82-1.60 (m, 2H).

EXAMPLE 5



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydropyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

15 A. 4-(1H-Pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

A solution of 1.9 g (8.47 mmol) of 1H-pyrrolo[3,2-c]pyridine (synthesized following the procedure described in *Synthesis*, 1996, 882), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (3.4 g, 16.9 mmol) and potassium hydroxide (1.9 g, 33.9 mmol) in MeOH (20 mL) was heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (100 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 50 mL). The organic extracts was dried over Na₂SO₄ and concentrated to form a solid. The solid was washed with MeOH (50 mL), filtered and dried to give a light yellow solid 2.0 g (79%). TLC (silica, 10% MeOH/CH₂Cl₂): $R_f = 0.5$. MS (electrospray): exact mass calculated for C₁₇H₂₁N₃O₂, 299.16; m/z found, 300.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 12.26 (br s, 1H), 9.20 (s, 1H), 8.28 (d, $J = 5.67$ Hz, 1H), 7.35 (dd, $J = 5.09$ Hz, 0.78 Hz, 1H), 7.32 (s, 1H), 6.19 (br s, 1H), 4.14 (br s, 2H), 3.68 (t, $J = 5.67$ Hz, 2H), 2.61-2.55 (m, 2H), 1.48 (s, 9H).

B. 4-(1H-Pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

4-(1H-Pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (2 g, 6.6 mmol) in EtOH (50 mL) containing PtO₂ (500 mg) was

placed on a Parr hydrogenator at 60 psi H₂. After 18 h the mixture was filtered through celite and evaporated to give a white solid (2.0 g, 100%). TLC (silica,

10% MeOH/CH₂Cl₂): R_f = 0.49. MS (electrospray): exact mass calculated for C₁₇H₂₃N₃O₂, 301.18; *m/z* found, 302.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz):

13.66 (br s, 1H), 8.88 (s, 1H), 8.79 (d, *J* = 6.46 Hz, 1H), 7.69 (d, *J* = 6.46 Hz,

1H), 7.30 (s, 1H), 4.14 (br s, 2H), 2.99-2.87 (m, 1H), 2.86-2.71 (m, 2H), 1.91 (d, *J* = 11.54 Hz, 2H), 1.64-1.50 (m, 2H), 1.38 (s, 9H).

C. 3-Piperidin-4-yl-1H-pyrrolo[3,2-c]pyridine.

4-(1H-Pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (2.0

g, 6.6 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated and the golden oil brought up in Et₂O. A solid formed and was

filtered, washed with Et₂O and air dried to give 2.1 g (100%) of a white solid as a TFA salt. MS (electrospray): exact mass calculated for C₁₂H₁₅N₃, 201.13; *m/z*

found, 202.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 9.4 (br s, 1H), 8.96 (s, 1H),

8.26 (d, *J* = 5.87 Hz, 1H), 7.24 (s, 1H), 6.99 (s, 1H), 3.22-3.16 (m, 2H), 3.05-

2.95 (m, 1H), 2.86-2.77 (m, 2H), 2.05 (d, *J* = 12.72 Hz, 2H), 1.89 (br s, 1H),

1.75-1.63 (m, 2H).

D. 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-

pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-
propan-2-ol.

3-Piperidin-4-yl-1H-pyrrolo[3,2-c]pyridine (159 mg, 0.5 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-

tetrahydro-1H-pyrazolo[4,3-c]pyridine (200 mg, 0.5 mmol) were set stirring in


EtOH (10 mL) containing Et₃N (112 μL, 0.77 mmol) at 80 °C. After 16 h the mixture was cooled, evaporated, brought up in CH₂Cl₂ and washed with water.

The organics were dried over Na₂SO₄ and concentrated. Column

chromatography (silica, 0-10% (2 N NH₃ in MeOH)/CH₂Cl₂) provided 82 mg

(27%) of a white solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.8. MS (electrospray): exact mass calculated for C₂₉H₃₃F₃N₆O₃S, 602.23; *m/z* found, 603.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 9.62 (s, 1H), 8.90 (s, 1H), 8.21 (d, *J* = 5.87 Hz, 1H), 7.69 (d, *J* = 7.83 Hz, 2H), 7.62 (d, *J* = 8.41 Hz, 2H), 7.23 (d, *J* = 5.87, 1H), 6.97 (s, 1H), 4.51 (dd, *J* = 14.48 Hz, 8.80 Hz, 2H), 4.23-4.13 (m, 2H), 4.05-3.95 (m, 1H), 3.72-3.54 (m, 3H), 3.11-2.98 (m, 2H), 2.95-2.86 (m, 2H), 2.84 (s, 3H), 2.51-2.39 (m, 3H), 2.20-2.11 (m, 1H), 2.07-1.97 (m, 2H), 1.85-1.63 (m, 2H).

10 EXAMPLE 6



15 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

A. 4-(1H-Pyrrolo[2,3-b]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

1H-Pyrrolo[2,3-b]pyridine (3 g, 25 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (4.2 g, 21 mmol) and potassium hydroxide (3.56 g, 63 mmol) were added in MeOH (60 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (300 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 150 mL). The organic extracts were dried over Na₂SO₄ and concentrated to form a solid. The solid was washed with MeOH (150 mL), filtered and dried to give a light yellow solid 5.7 g (91%). TLC (silica, 50% EtOAc/hexanes): R_f = 0.3. MS (electrospray): exact mass calculated for C₁₇H₂₁N₃O₂, 299.16; *m/z* found, 300.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz) 10.97 (br s, 1H), 8.33 (dd, *J* = 3.33 Hz, 1.37 Hz, 1H), 8.20 (dd, *J* = 6.65 Hz, 1.37 Hz, 1H), 7.34 (br s, 1H), 7.25 (s, 1H),

7.13 (dd, $J = 4.89$ Hz, 3.13 Hz, 1H), 4.14 (br s, 2H), 3.68 (t, $J = 5.28$ Hz, 2H), 2.56 (br s, 2H), 1.49 (s, 9H).

B. 4-(1H-Pyrrolo[2,3-b]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

5 4-(1H-Pyrrolo[2,3-b]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (1 g, 3.3 mmol) in EtOH (25 mL) containing PtO₂ (250 mg) was placed on a Parr hydrogenator at 60 psi H₂. After 18 h the mixture was filtered through celite and evaporated to give 0.96 g (97%) of a white solid. TLC (silica, 50% EtOAc/hexanes): $R_f = 0.5$. MS (electrospray): exact mass
10 calculated for C₁₇H₂₃N₃O₂, 301.18; m/z found, 302.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz) 10.95 (br s, 1H), 8.26 (dd, $J = 3.33$ Hz, 1.37 Hz, 1H), 7.96 (dd, $J = 6.26$ Hz, 1.57 Hz, 1H), 7.11 (s, 1H), 7.05 (dd, $J = 4.89$ Hz, 3.13 Hz, 1H), 4.22 (br s, 2H), 3.00-2.79 (m, 3H), 1.99 (d, $J = 13.89$ Hz, 2H), 1.74-1.60 (m, 2H), 1.47 (s, 9H).

15 C. 3-Piperidin-4-yl-1H-pyrrolo[2,3-b]pyridine.


4-(1H-Pyrrolo[2,3-b]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (963 mg, 3.2 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated and the golden oil brought up in Et₂O. A solid formed
20 and was filtered, washed with Et₂O and air dried to give 974 mg (96%) of a white solid as a TFA salt. MS (electrospray): exact mass calculated for C₁₂H₁₅N₃, 201.13; m/z found, 202.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.09 (dd, $J = 3.33$ Hz, 1.57 Hz, 1H), 7.89 (dd, $J = 6.26$ Hz, 1.57 Hz, 1H), 7.01 (s, 1H), 6.99 (dd, $J = 4.89$ Hz, 3.13 Hz, 1H), 5.04 (br s, 2H), 3.11-3.04 (m, 2H),
25 2.88-2.79 (m, 1H), 2.73-2.64 (m, 2H), 1.94 (d, $J = 12.52$ Hz, 2H), 1.65-1.63 (m, 2H).

D. 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

30 3-Piperidin-4-yl-1H-pyrrolo[2,3-b]pyridine (443 mg, 1.4 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (289 mg, 0.7 mmol) were set stirring in

EtOH (10 mL) containing Et₃N (146 μ L, mmol) at 80 °C. After 16 h the mixture was cooled, evaporated, brought up in CH₂Cl₂ and washed with water. The organics were dried over Na₂SO₄ and concentrated. Column chromatography (silica, 0-10% (2 N NH₃ in MeOH)/CH₂Cl₂) provided 107 mg (25%) of a white solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.45. MS (electrospray): exact mass calculated for C₂₉H₃₃F₃N₆O₃S, 602.23; *m/z* found, 603.3 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 10.6 (br s, 1H), 8.24 (m, 1H), 7.91 (m, 1H), 7.70 (m, 2H), 7.63 (m, 2H), 7.05 (br s, 1H), 7.02 (m, 1H), 4.52 (dd, *J* = 4.28 Hz, 9.78 Hz, 2H), 4.24-4.16 (m, 2H), 4.06-3.98 (m, 1H), 3.72-3.64 (m, 1H), 3.64-3.55 (m, 1H), 3.11-3.00 (m, 2H), 2.96-2.87 (m, 2H), 2.85 (s, 3H), 2.82-2.74 (m, 1H), 2.53-2.40 (m, 3H), 2.22-2.12 (m, 1H), 2.05-1.95 (m, 2H), 1.85-1.64 (m, 2H).

EXAMPLE 7



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydropyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

A. [2-(2-Chloro-5-nitro-pyridin-4-yl)-vinyl]-dimethyl-amine.

A solution of 2-chloro-4-methyl-5-nitro-pyridine (2 g, 11.59 mmol) in DMF (11.6 mL) was treated with 3.08 mL (23.2 mmol, 2 eq) of DMF-dimethylacetal and the reaction mixture was stirred at 100 °C for 4 h. All volatiles were removed under reduced pressure. Column chromatography (silica, 20% EtOAc/hexanes) provided 2.37 g (90%) of [2-(2-chloro-5-nitro-pyridin-4-yl)-vinyl]-dimethyl-amine. TLC (silica, 20% EtOAc/hexanes): R_f = 0.30. MS (electrospray): exact mass calculated for C₉H₁₀ClN₃O₂, 227.05; *m/z* found, 228.1 [M+H]⁺. ¹H NMR (400

MHz, CDCl₃): 8.79 (s, 1H), 8.02 (s, 1H), 7.35 (d, *J* = 13 Hz, 1H), 5.94 (d, *J* = 13 Hz, 1H), 2.96 (s, 3H), 2.87 (s, 3H).

B. 5-Morpholin-4-yl-1H-pyrrolo[2,3-c]pyridine.

- 5 A solution of 450 mg (2 mmol) of [2-(2-chloro-5-nitro-pyridin-4-yl)-vinyl]-dimethyl-amine in a 20 mL of mixed solvent of MeOH-CH₂Cl₂ (1:1) was treated with 3 mL of morpholine. The reaction mixture was stirred at 65 °C for 8 h. Volatiles were then removed. CH₂Cl₂ (100 mL) and H₂O (30 mL) were added. The organic layer was separated and washed with H₂O (30 mL), brine (30 mL),
10 dried over Na₂SO₄, and concentrated. The red powder was treated with 4.0 g (63 mmol, 32 eq) of ammonium formate and 10% Pd-C (120 mg). The reaction mixture was stirred at 65 °C for 30 min. The reaction mixture was then filtered through a pad of celite and concentrated to obtain a yellow solid. Column chromatography (silica, 5% MeOH/CH₂Cl₂) provided 210 mg (52% for 2 steps)
15 of 5-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridine as a yellow solid. TLC (silica, 5% MeOH/CH₂Cl₂): *R_f* = 0.40. MS (electrospray): exact mass calculated for C₁₁H₁₃N₃O, 203.11; *m/z* found, 204.2 [M+H]⁺.


C. 4-(5-Morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

- 20 A solution of 200 mg (1.0 mmol) of 5-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridine and 398 mg (2.0 mmol, 2 eq) of 4-oxo-piperidine-1-carboxylic acid tert-butyl ester in 5 mL of MeOH was treated with 224 mg (4.0 mmol, 4 eq) of potassium hydroxide. The reaction mixture was stirred at 65 °C for 12 h and volatiles were
25 removed. The crude product was partitioned between CH₂Cl₂ (100 mL) and 20 mL of H₂O. The organic layer was washed with H₂O (2 x 20 mL), dried over Na₂SO₄ and concentrated. The yellow powder was treated with 630 mg (10 mmol, 10 eq) of ammonium formate and 10% Pd-C (50 mg). The reaction mixture was stirred at 65 °C for 1 h. The reaction mixture was then filtered
30 through a pad of celite and concentrated to obtain a yellow solid. Column chromatography (silica, 5% MeOH/CH₂Cl₂) provided 180 mg (47% for 2 steps) of a yellow solid. TLC (silica, 5% MeOH/CH₂Cl₂): *R_f* = 0.40. MS (electrospray): exact mass calculated for C₂₁H₃₀N₄O₃, 386.23; *m/z* found, 387.2 [M+H]⁺.

D. 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydropyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

- 5 4-(5-Morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (180 mg, 0.47 mmol) was dissolved in 3.0 mL of CH₂Cl₂ and treated with 2.5 mL of trifluoroacetic acid. The reaction mixture was stirred at 25 °C for 1 h before all volatiles were removed. The solid was dissolved in MeOH (20 mL) and neutralized with DOWEX 550A OH anion exchange resin
- 10 to pH 8. The resin was then filtered off and MeOH was removed under reduced pressure. The residue was dissolved in 2.5 mL of iPrOH and treated with 187 mg (0.47 mmol, 1 eq) of 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine. The reaction was stirred at 85 °C for 3 h before solvent was removed. Column
- 15 chromatography (silica, 5-10% MeOH/CH₂Cl₂ then 5-10% (2 N NH₃ in MeOH)/CH₂Cl₂) provided 97 mg (30%) of the title compound. TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.25. MS (electrospray): exact mass calculated for C₃₃H₄₀F₃N₇O₄S, 687.28; *m/z* found, 688.3 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 8.47 (br s, 1H), 8.44 (d, *J* = 1.0 Hz, 1H), 7.70 and 7.65 (AB pattern, *J* = 8.4 Hz, 4H), 7.03 (d, *J* = 2.1 Hz, 1H), 6.76 (s, 1H), 4.58- 4.50 (m, 2H), 4.21-4.00 (m, 3H), 3.90 (t, *J* = 4.5 Hz, 4H), 3.72-3.58 (m, 2H), 3.40 (t, *J* = 4.5 Hz, 4H), 3.10-2.85 (m, 4H), 2.88 (s, 3H), 2.80-2.70 (m, 1H), 2.52-2.41 (m, 3H), 2.20-2.00 (m, 3H), 1.80-1.60 (m, 2H).
- 20

EXAMPLE 8



1-[4-(6-Dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

A. Dimethyl-(5-methyl-4-nitro-1-oxy-pyridin-2-yl)-amine.

A solution of 2-bromo-5-methyl-4-nitro-pyridine 1-oxide (674 mg, 2.78 mmol) in 2 M dimethylamine in methanol (20 mL) was heated at 65 °C for 16 h. The solvent was evaporated under reduced pressure. The residue was purified by column chromatography (silica, 30-80% EtOAc/hexanes) to obtain 290 mg (53%) of the desired product. TLC (silica, 50% EtOAc/hexanes): R_f = 0.10. MS (electrospray): exact mass calculated for $C_8H_{11}N_3O_3$, 197.08; m/z found, 198.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.04 (s, 1H), 7.50 (s, 1H), 3.00 (s, 6H), 2.44 (s, 3H).

B. Dimethyl-(1H-pyrrolo[3,2-c]pyridin-6-yl)-amine.

A solution of dimethyl-(5-methyl-4-nitro-1-oxy-pyridin-2-yl)-amine (290 mg, 1.47 mmol) in DMF (3 mL) was treated with DMF-dimethylacetal (390 μ L, 2.94 mmol) and the reaction mixture was stirred at 100 °C for 4 h. All volatiles were removed under reduced pressure. The red powder was treated with ammonium formate (927 mg, 14.7 mmol) and 10% Pd-C (156 mg). The reaction mixture was stirred at 65 °C for 30 min. The reaction mixture was then filtered through a pad of celite and concentrated to obtain a yellow solid. Column chromatography (silica, 5% MeOH/CH₂Cl₂) provided 100 mg (42% for two steps) of product as a yellow solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.2. MS (electrospray): exact mass calculated for $C_9H_{11}N_3$, 161.10; m/z found,

162.1 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.55 (s, 1H), 8.28 (br s, 1H), 6.96 (dd, *J* = 1.96 Hz, 1.37 Hz, 1H), 6.45-6.43 (m, 1H), 6.39 (s, 1H), 3.08 (s, 6H).

C. 4-(6-Dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

Dimethyl-(1H-pyrrolo[3,2-c]pyridin-6-yl)-amine (100 mg, 0.62 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (248 mg, 1.24 mmol) and potassium hydroxide (139 mg, 2.48 mmol) were added in MeOH (5 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (20 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 10 mL). The organic extracts were dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography (silica, 0-5% MeOH/CH₂Cl₂) to obtain 180 mg (85%) of the title compound. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.58. MS (electrospray): exact mass calculated for C₁₉H₂₆N₄O₂, 342.21; *m/z* found, 343.2 (M⁺+H). ¹H NMR (CDCl₃, 400 MHz): 9.09 (br s, 1H), 8.76 (s, 1H), 6.90 (d, *J* = 2.15 Hz, 1H), 6.32 (s, 1H), 6.10 (br s, 1H), 4.10-4.05 (m, 2H), 3.62 (t, *J* = 5.87 Hz, 2H), 3.03 (s, 6H), 2.52-2.44 (m, 2H), 1.46 (s, 9H).

D. 4-(6-Dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

A solution of 4-(6-dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (180 mg, 0.53 mmol) in MeOH (10 mL) was treated with ammonium formate (332 mg, 5.3 mmol) and 10% Pd-C (56 mg). The reaction mixture was stirred at 65 °C for 1 h. The reaction mixture was then filtered through a pad of celite and concentrated to obtain an off white solid. Column chromatography (silica, 0-5% MeOH/CH₂Cl₂) provided 135 mg (75%) of product as a white solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.50. MS (electrospray): exact mass calculated for C₁₉H₂₈N₄O₂, 344.22; *m/z* found, 345.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.53 (s, 1H), 8.30 (br s, 1H), 6.67 (dd, *J* = 1.17 Hz, 0.78 Hz, 1H), 6.34 (d, *J* = 0.78 Hz, 1H), 4.33-4.16 (m, 2H), 3.05 (s, 6H), 2.97-2.80 (m, 3H), 1.99 (d, *J* = 12.72 Hz, 2H), 1.69-1.53 (m, 2H), 1.46 (s, 9H).

E. Dimethyl-(3-piperidin-4-yl-1H-pyrrolo[3,2-c]pyridin-6-yl)-amine.


The 4-(6-dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (135 mg, 0.39 mmol) was set stirring in 1:1 TFA/CH₂Cl₂.

- 5 After 45 min the mixture was evaporated. The residue was dissolved in MeOH (10 mL) and neutralized with DOWEX 550A OH anion exchange resin to pH 8. The resin was then filtered off and MeOH was removed under reduced pressure to give 96 mg (100%) of a yellow solid. MS (electrospray): exact mass calculated for C₁₄H₂₀N₄, 244.17; *m/z* found, 245.2 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 9.23 (br s, 1H), 8.54 (s, 1H), 6.67 (s, 1H), 6.31 (d, *J* = 0.98 Hz, 1H), 3.15 (d, *J* = 2.13 Hz, 2H), 3.02 (s, 6H), 2.91-2.81 (m, 1H), 2.81-2.72 (m, 2H), 2.01 (d, *J* = 12.52 Hz, 2H), 1.69-1.52 (m, 2H).

F. 1-[4-(6-Dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

- 15 Dimethyl-(3-piperidin-4-yl-1H-pyrrolo[3,2-c]pyridin-6-yl)-amine (96 mg, 0.53 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (319 mg, 0.80 mmol) were set stirring in ¹PrOH (10 mL) at 80 °C. After 16 h the mixture was cooled and concentrated. The residue was purified by column chromatography (silica, 0-10% (2 N NH₃ in MeOH)/CH₂Cl₂ to obtain 61 mg (18%) of a white solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.12. MS (electrospray): exact mass calculated for C₃₁H₃₈F₇N₇O₃S, 645.27; *m/z* found, 646.3 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.53 (s, 1H), 7.93 (br s, 1H), 7.71 (d, *J* = 8.22 Hz, 2H), 7.64 (d, *J* = 8.22, 2H), 6.67 (br s, 1H), 6.33 (d, *J* = 0.98 Hz, 1H), 4.54 (dd, *J* = 14.28 Hz, 9.59 Hz, 2H), 4.22-4.10 (m, 2H), 4.04-3.97 (m, 1H), 3.74-3.57 (m, 2H), 3.13-3.06 (m, 1H), 3.05 (s, 6H), 3.03-2.87 (m, 3H), 2.85 (s, 3H), 2.82-2.71 (m, 1H), 2.50-2.37 (m, 3H), 2.20-2.11 (m, 1H), 2.06-1.97 (m, 2H), 1.82-1.61 (m, 2H).
- 20
25
30

EXAMPLE 9



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-
 5 pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-
 piperidin-1-yl]-propan-2-ol.

A. 4-(5-Methyl-4-nitro-1-oxy-pyridin-2-yl)-morpholine.

A solution of 2-bromo-5-methyl-4-nitro-pyridine 1-oxide (500 mg, 2.14 mmol) in
 10 morpholine (15 mL) was heated at 70 °C for 16 h. The solvent was evaporated
 under reduced pressure. The residue was purified by column chromatography
 (silica, 30-80% EtOAc/hexanes) to obtain 480 mg (94%) of the desired product.
 TLC (silica, 50% EtOAc/hexanes): R_f = 0.10. MS (electrospray): exact mass
 calculated for $C_{10}H_{13}N_3O_4$, 239.09; m/z found, 240.1 [$M^+ + H$]. 1H NMR ($CDCl_3$,
 15 400 MHz): 8.09 (s, 1H), 7.55 (s, 1H), 3.90 (t, J = 4.50 Hz, 4H), 3.36 (t, J = 4.70
 Hz, 4H), 2.50 (s, 3H).

B. 6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridine.

A solution of 4-(5-methyl-4-nitro-1-oxy-pyridin-2-yl)-morpholine (480 mg, 2
 20 mmol) in DMF (5 mL) was treated with DMF-dimethylacetal (533 μ L, 4 mmol)
 and the reaction mixture was stirred at 100 °C for 4 h. All volatiles were
 removed under reduced pressure. The red powder was treated with
 ammonium formate (1.26 g, 20 mmol) and 10% Pd-C (212 mg). The reaction
 mixture was stirred at 65 °C for 30 min. The reaction mixture was then filtered
 25 through a pad of celite and concentrated to obtain a yellow solid. Column
 chromatography (silica, 5% MeOH/ CH_2Cl_2) provided 197 mg (49% for two
 steps) of a yellow solid. TLC (silica, 10% MeOH/ CH_2Cl_2): R_f = 0.55. MS

(electrospray): exact mass calculated for $C_{11}H_{13}N_3O$, 203.11; m/z found, 204.1 $[M+H]^+$. 1H NMR ($CDCl_3$, 400 MHz) 8.68 (br s, 1H), 8.59 (s, 1H), 7.04 (dd, $J = 2.15$ Hz, 1.17 Hz, 1H), 6.51 (s, 1H), 6.49-6.47 (m, 1H), 3.86 (t, $J = 4.70$ Hz, 4H), 3.40 (t, $J = 4.70$ Hz, 4H).

5

C. 4-(6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridine (197 mg, 0.97 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (387 mg, 1.94 mmol) and potassium hydroxide (218 mg, 3.88 mmol) were added in MeOH (10 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and poured into ice water (50 mL). The mixture was extracted with 10% MeOH/ CH_2Cl_2 (5 x 20 mL). The organic extracts were dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography (silica, 0-5% MeOH/ CH_2Cl_2) to obtain 337 mg (91%) of the desired product. TLC (silica, 5% MeOH/ CH_2Cl_2): $R_f = 0.50$. MS (electrospray): exact mass calculated for $C_{21}H_{28}N_4O_3$, 384.22; m/z found, 749.3 $[M^+ + H]$. 1H NMR ($CDCl_3$, 400 MHz): 9.24 (br s, 1H), 8.77 (s, 1H), 6.95 (d, $J = 2.15$ Hz, 1H), 6.45 (s, 1H), 6.09 (br s, 1H), 4.09-4.04 (m, 2H), 3.66 (t, $J = 6.06$ Hz, 4H), 3.60 (t, $J = 5.28$ Hz, 2H), 3.25 (t, $J = 5.09$ Hz, 4H), 2.38 (d, $J = 6.26$ Hz, 2H), 1.44 (s, 9H).

20

D. 4-(6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

4-(6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (337 mg, 0.9 mmol) in MeOH (20 mL) was treated with ammonium formate (568 mg, 9.0 mmol) and 10% Pd-C (95 mg). The reaction mixture was stirred at 65 °C for 1 h. The reaction mixture was then filtered through a pad of celite and concentrated to obtain an off-white solid. Column chromatography (silica, 0-5% MeOH/ CH_2Cl_2) provided 340 mg (98%) of a white solid. TLC (silica, 5% MeOH/ CH_2Cl_2): $R_f = 0.40$. MS (electrospray): exact mass calculated for $C_{21}H_{30}N_4O_3$, 386.23; m/z found, 387.3 $[M^+ + H]$. 1H NMR ($CDCl_3$, 400 MHz) 9.14 (br s, 1H), 8.55 (s, 1H), 6.74 (d, $J = 1.96$ Hz, 1H), 6.45 (s, 1H), 4.27-4.08 (m, 2H), 3.80 (t, $J = 4.50$ Hz, 4H), 3.34 (t,

30

$J = 4.89$ Hz, 4H), 2.96-2.77 (m, 3H), 1.97 (d, $J = 12.91$ Hz, 2H), 1.67-1.52 (m, 2H), 1.44 (s, 9H).


E. 6-Morpholin-4-yl-3-piperidin-4-yl-1H-pyrrolo[3,2-c]pyridine.

- 5 4-(6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (380 mg, 0.98 mmol) was set stirring in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated. The residue was dissolved in MeOH (20 mL) and neutralized with DOWEX 550A OH anion exchange resin to pH 8. The resin was then filtered off and MeOH was removed under reduced
- 10 pressure to give 281 mg (100%) of a yellow solid. MS (electrospray): exact mass calculated for C₁₆H₂₂N₄O, 286.18; m/z found, 287.1 [M⁺+H].

F. 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

- 15 6-Morpholin-4-yl-3-piperidin-4-yl-1H-pyrrolo[3,2-c]pyridine (281 mg, 0.98 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (591 mg, 1.47 mmol) were set stirring in ⁱPrOH (10 mL) at 80 °C. After 16 h the mixture was cooled and concentrated.
- 20 The residue was purified by column chromatography (silica, 0-10% (2 *N* NH₃ in MeOH)/CH₂Cl₂) to obtain 468 mg (69%) of a white solid. TLC (silica, 10% (2 *N* NH₃ in MeOH)/CH₂Cl₂): $R_f = 0.62$. MS (electrospray): exact mass calculated for C₃₃H₄₀F₃N₇O₄S, 687.28; m/z found, 688.3 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.56 (s, 1H), 8.40 (br s, 1H), 7.69 (d, $J = 8.41$ Hz, 2H), 7.63 (d, $J = 8.41$, 2H), 6.73 (br s, 1H), 6.46 (s, 1H), 4.51 (dd, $J = 14.28$ Hz, 8.80 Hz, 2H), 4.21-4.10
- 25 (m, 2H), 4.03-3.95 (m, 1H), 3.82 (t, $J = 4.11$ Hz, 4H), 3.71-3.54 (m, 2H), 3.36 (t, $J = 4.89$ Hz, 4H), 3.10-2.97 (m, 2H), 2.93-2.86 (m, 2H), 2.84 (s, 3H), 2.82-2.72 (m, 1H), 2.50-2.37 (m, 3H), 2.20-2.10 (m, 1H), 2.04-1.95 (m, 2H), 1.80-1.60 (m, 2H).

EXAMPLE 10



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

A. 6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridine 5-oxide.

A solution of 4-(5-methyl-4-nitro-1-oxy-pyridin-2-yl)-morpholine (480 mg, 2 mmol) in DMF (5 mL) was treated with DMF-dimethylacetal (533 μ L, 4 mmol) and the reaction mixture was stirred at 100 °C for 4 h. All volatiles were removed under reduced pressure. The red powder was treated with ammonium formate (1.26 g, 20 mmol) and 10% Pd-C (212 mg). The reaction mixture was stirred at 65 °C for 30 min. The reaction mixture was then filtered through a pad of celite and concentrated to obtain a yellow solid. Column chromatography (silica, 5% MeOH/CH₂Cl₂) provided 130 mg (30% for two steps) of a yellow solid. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.28. MS (electrospray): exact mass calculated for C₁₁H₁₃N₃O₂, 219.10; *m/z* found, 220.1 [M+H]⁺. ¹H NMR (CDCl₃, 400 MHz): 8.42 (br s, 1H), 7.25 (s, 1H), 7.14 (s, 1H), 6.82 (s, 1H), 6.35 (s, 1H), 3.79 (t, *J* = 4.70 Hz, 4H), 3.18 (t, *J* = 4.70 Hz, 4H).

B. 4-(6-Morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester.

6-Morpholin-4-yl-1H-pyrrolo[3,2-c]pyridine 5-oxide (130 mg, 0.59 mmol), 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (237 mg, 1.19 mmol) and potassium hydroxide (133 mg, 2.37 mmol) were added in MeOH (8 mL) and heated to reflux for 16 h. The reaction mixture was then cooled to room temperature and

poured into ice water (30 mL). The mixture was extracted with 10% MeOH/CH₂Cl₂ (5 x 10 mL). The organic extracts were dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography (silica, 0-5% MeOH/CH₂Cl₂) to obtain 140 mg (59%) of the desired product. TLC (silica, 10% MeOH/CH₂Cl₂): R_f = 0.55. MS (electrospray): exact mass calculated for C₂₁H₂₈N₄O₄, 400.21; *m/z* found, 401.2 (M⁺+H).

C. 4-(6-Morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

4-(6-Morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (140 mg, 0.35 mmol) in EtOH (20 mL) containing PtO₂ (50 mg) was placed on a Parr hydrogenator at 60 psi H₂. After 18 h mixture was filtered through celite and evaporated to give a white solid. Column chromatography (silica, 0-5% (2 N NH₃ in MeOH)/CH₂Cl₂) provided 56 mg (40%) of a white solid. TLC (silica, 5% MeOH/CH₂Cl₂): R_f = 0.13. MS (electrospray): exact mass calculated for C₂₁H₃₀N₄O₄, 402.23; *m/z* found, 403.2 (M⁺+H). ¹H NMR (CDCl₃, 400 MHz): 11.63 (br s, 1H), 8.58 (s, 1H), 6.97 (br s, 1H), 6.69 (s, 1H), 4.26-4.08 (m, 2H), 3.73 (t, *J* = 4.30 Hz, 4H), 3.17 (t, *J* = 4.50 Hz, 4H), 2.92-2.74 (m, 3H), 1.91 (d, *J* = 11.93 Hz, 2H), 1.62-1.50 (m, 2H), 1.43 (s, 9H).


D. 6-Morpholin-4-yl-3-piperidin-4-yl-1H-pyrrolo[3,2-c]pyridine 5-oxide.

4-(6-Morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (56 mg, 0.14 mmol) was stirred in 1:1 TFA/CH₂Cl₂. After 45 min the mixture was evaporated. The residue was dissolved in MeOH (10 mL) and neutralized with DOWEX 550A OH anion exchange resin to pH 8. The resin was then filtered off and MeOH was removed under reduced pressure to give 42 mg (100%) of a yellow solid. MS (electrospray): exact mass calculated for C₁₆H₂₂N₄O₂, 302.17; *m/z* found, 303.1 [M⁺+H].

E. 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

6-Morpholin-4-yl-3-piperidin-4-yl-1H-pyrrolo[3,2-c]pyridine 5-oxide (42 mg, 0.14 mmol) and 5-methanesulfonyl-1-oxiranylmethyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (84 mg, 0.21 mmol) were set stirring in ⁱPrOH (5 mL) at 80 °C. After 6 h the mixture was cooled and concentrated. The residue was purified by column chromatography (silica, 0-10% (2 N NH₃ in MeOH)/CH₂Cl₂) to obtain 5.1 mg (5%) of a white solid. TLC (silica, 10% (2 N NH₃ in MeOH)/CH₂Cl₂): R_f = 0.54. MS (electrospray): exact mass calculated for C₃₃H₄₀F₃N₇O₅S, 703.28; *m/z* found, 704.3 [M⁺+H]. ¹H NMR (CDCl₃, 400 MHz): 8.60 (s, 1H), 7.71 (d, *J* = 8.41 Hz, 2H), 7.66 (d, *J* = 8.41, 2H), 7.27 (s, 1H), 6.95 (s, 1H), 6.70 (s, 1H), 4.55 (dd, *J* = 14.48 Hz, 3.13 Hz, 2H), 4.23-4.11 (m, 2H), 4.05-3.97 (m, 1H), 3.84 (t, *J* = 4.30 Hz, 4H), 3.72-3.60 (m, 2H), 3.23 (t, *J* = 4.30 Hz, 4H), 3.12-2.98 (m, 2H), 2.97-2.89 (m, 2H), 2.88 (s, 3H), 2.73-2.63 (m, 1H), 2.51-2.36 (m, 3H), 2.17-2.08 (m, 1H), 1.99-1.90 (m, 2H), 1.78-1.58 (m, 2H).

EXAMPLE 11



6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-hydroxy-ethyl)-amide.

A. 1-[4-(2,4-Difluoro-benzoyl)-piperidin-1-yl]-ethanone.

A stirred solution of 10 g (58.5 mmol) of 1-acetylpiperidine-4-carboxylic acid in anhydrous dichloroethane (35 mL) was treated with 5.1 mL (70.2 mmol) of thionyl chloride in 7 mL of dichloroethane and then heated to 60 °C for 30 min. Another flask containing a suspension of 8.02 mL (81.8 mmol) of 1,3-difluorobenzene and 17.9 g (134 mmol) of aluminum chloride in 55 mL of dichloroethane was prepared, to this was added the previously prepared acid

chloride suspension in portions. The resulting suspension was refluxed for 4 h, cooled and then poured over ice and HCl. The acidic solution was extracted with CH₂Cl₂ (3 x 300 mL) and the combined organic extracts were washed with brine, dried over Na₂SO₄, and concentrated. The crude product was

5 recrystallized from hexanes to afford 9.5 g (61%) of the desired product as a white solid. MS (electrospray): exact mass calculated for C₁₄H₁₅F₂NO₂, 267.11; m/z found, 268.1 [M+H]⁺, ¹H NMR (CDCl₃, 400 MHz): 7.87 (dt, J = 8.41, 6.65 Hz 1H), 6.98 (m, 1H), 6.88 (ddd, J = 10.96, 8.61, 2.35 Hz, 1H), 4.55 (m, 1H), 3.87 (m, 1H), 3.32 (dtt, J = 10.76, 3.91, 1.37 Hz, 1H), 3.19 (ddd, J = 13.89, 11.93, 2.93 Hz, 1H), 2.79 (ddd, J = 13.89, 12.24, 2.93 Hz, 1H), 2.10 (s, 3H), 1.95 (br d, J = 12.91 Hz, 2H), 1.72 (br m, 1H), 1.56 (br m, 1H).

B. 3-(1-Acetyl-piperidin-4-yl)-6-fluoro-benzo[b]thiophene-2-carboxylic acid methyl ester.

15 To a stirred solution of 33.6 g (0.126 mol) of 1-[4-(2,4-difluoro-benzoyl)-piperidin-1-yl]-ethanone and 13 mL (145 mol) of methyl thioglycolate in 320 mL dry THF was added 5.8 g (145 mol) of 60% sodium hydride in mineral oil in portions. The reaction mixture was heated to reflux overnight, allowed to cool to room temperature and the solvent removed under reduced pressure. The residue was then partitioned between 300 mL of CH₂Cl₂ and 200 mL of water. The aqueous layer was further extracted with CH₂Cl₂ (2 x 500 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated to give a residue which was then triturated with hexanes/EtOAc to give 27.5 g (65%) of the desired product as a white solid. MS (electrospray):
25 exact mass calculated for C₁₇H₁₈FNO₃S, 335.1; m/z found, 336.1 [M+H]⁺. ¹H NMR (DMSO-d₆, 400 MHz, a mixture of amide rotamers): 7.12 (m, 2H), 6.92 (dt, J = 8.41, 1.77 Hz, 1H), 4.43 (d, J = 3.79 Hz, 1H), 4.43-4.36 (m, 1H), 3.82 (bt, J = 14.65 Hz, 1H), 3.57 (s, 3H), 2.92-2.79 (m, 1H), 2.38-2.34 (m, 1H), 1.94 (s, 1.5H), 1.93 (s, 1.5H), 1.86-1.72 (m, 1H), 1.47-1.38 (m, 1H), 1.38-1.27 (m, 0.5H), 1.27-1.16 (m, 1H), 1.15-1.03 (m, 0.5H).
30

C. 6-Fluoro-3-piperidin-4-yl-benzo[b]thiophene-2-carboxylic acid methyl ester hydrochloride salt.

A solution of 24.4 g (66.8 mmol) of 3-(1-acetyl-piperidin-4-yl)-6-fluoro-benzo[b]thiophene-2-carboxylic acid methyl ester in 400 mL of MeOH and 50 mL of concentrated HCl was heated a reflux for 2 days. When the solution was allowed to cool to room temperature the white precipitate was filtered, washed with methanol and dried to give 17.9 g (74%) of product as a white powder.

MS (electrospray): exact mass calculated for $C_{15}H_{16}FNO_2S$, 293.09; m/z found, 294.1 $[M+H]^+$. 1H NMR (DMSO- d_6 , 400 MHz): 9.38 (br s, 1H), 9.02 (br s, 1H), 8.60 (dd, $J = 9.19, 5.09$ Hz, 1H), 7.98 (dd, $J = 9.00, 2.54$ Hz, 1H), 7.36 (dt, $J = 9.00, 2.54$ Hz, 1H), 4.37 (br t, $J = 12.72$ Hz, 1H), 3.87 (s, 3H), 3.40 (br d, $J = 11.93$ Hz, 2H), 3.02 (q, $J = 11.35$ Hz, 2H), 2.61 (dq, $J = 13.30, 3.72$ Hz, 2H), 1.77 (br d, $J = 12.91$ Hz, 2H).

D. 3-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-1-ol.

Cs_2CO_3 (33.74 g, 103.5 mmol) was added to a solution of 5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine (29.8 g, 86.3 mmol) in anhydrous DMF (70 mL) and stirred for 25 min. 3-Bromo-1-propanol (8.6 mL, 13.2 g, 94.9 mmol) was added and stirred under N_2 at room temperature for 18 h. Water (500 mL) was added to the reaction and stirred for 5 min. The precipitated material was filtered out and washed with water (4 X 100 mL) and dried. The crude material (31.0 g) was taken up in anhydrous DMF (65 mL) and Cs_2CO_3 (33.74 g, 103.5 mmol) was added, and stirred for 10 min. Then 3-bromo-1-propanol (8.6 mL, 13.2 g, 94.9 mmol) and MeOH (6.0 mL, 4.75 g, 148 mmol) were added and stirring continued under N_2 at rt for 15 h. Water (500 mL) was added to the reaction and stirred for 10 min. The precipitated material was filtered and washed with water (3 X 100 mL). The filter cake was dissolved in CH_2Cl_2 (200 mL) and washed with brine (50 mL), dried (Na_2SO_4), and concentrated. The solid was triturated with Et_2O (200 mL), filtered, then washed with Et_2O , and dried to furnish 16.0 g of the desired compound. The mother liquor was chromatographed (silica, 0-10% acetone/EtOAc) to obtain an additional 3.0 g of the title compound. The combined yield was 54.6%. MS (electrospray): exact mass calculated for $C_{17}H_{20}F_3N_3O_3S$, 403.12; m/z found, 404.0 $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$):

7.71 (d, $J = 8.2$ Hz, 2H), 7.66 (d, $J = 8.5$ Hz, 2H), 4.55 (s, 2H), 4.23 (t, $J = 6.5$ Hz, 2H), 3.70-3.63 (m, 4H), 2.90 (s, 3H), 2.90 (t, $J = 5.1$ Hz, 2H), 2.62 (t, $J = 5.9$ Hz, 1H), 2.06 (q, $J = 6.1$ Hz, 2H).

5 E. 3-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propionaldehyde.

Dess-Martin periodinane (3.45 g, 8.2 mmol) was added to a solution of 3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-1-ol (3.0 g, 7.4 mmol) in CH_2Cl_2 (20 mL) at 0 °C under N_2 .

- 10 After 15 min, the reaction was allowed to warm to room temperature and stirred for another 1.5 h. The reaction was diluted with Et_2O (60 mL) and 20 % aq. NaHCO_3 (35 mL) was added slowly (caution! rapid gas evolution). Then $\text{Na}_2\text{S}_2\text{O}_3$ was added and stirred at room temperature for 30 min. The layers were separated and the aqueous portion was extracted with Et_2O (2x30 mL).
- 15 The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. MPLC (silica, 1-10% $\text{MeOH}/\text{CH}_2\text{Cl}_2$) afforded 2.53 g of the desired aldehyde in 85% yield. MS (electrospray): exact mass calculated for $\text{C}_{17}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_3\text{S}$, 401.11; m/z found, 402.1 $[\text{M}+\text{H}]$. ^1H NMR (400 MHz, CDCl_3): 9.82 (s, 1H), 7.63 (d, $J = 8.4$ Hz, 2H), 7.58 (d, $J = 8.4$ Hz, 2H), 4.68 (s, 2H), 4.25 (t, $J = 6.1$ Hz, 2H), 3.63 (t, $J = 5.8$ Hz, 4H), 3.14 (t, $J = 6.1$ Hz, 2H), 2.92 (t, $J = 5.8$ Hz, 2H), 2.81 (s, 3H).
- 20

25 F. 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid methyl ester.

- To a stirred solution of 410 mg (1.25 mmol) of 6-fluoro-3-piperidin-4-yl-benzo[b]thiophene-2-carboxylic acid methyl ester hydrochloride salt in 10 mL of dichloromethane and 0.18 mL (1.25 mmol) of triethylamine was added 500 mg of 3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propionaldehyde (1.25 mmol) and 2 g of NaHCO_3 .
- 30 The mixture was stirred for 4 h before the portion wise addition of 792 mg (3.73 mmol) sodium triacetoxyborohydride. The reaction was stirred at room temperature for 3 h before quenching with 20 mL of water. The aqueous

phase was extracted with CH₂Cl₂ (3 x 50 mL). The combined organic layers were then washed with brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (silica, 0-5% MeOH/CH₂Cl₂) to afford 650 mg (77%) of a white solid. MS (electrospray): exact mass calculated for C₃₂H₃₄F₄N₄O₄S₂: 678.20; *m/z* found, 679.2 [M+H]⁺. ¹H NMR (CDCl₃, 400 MHz): 7.74 and 7.66 (A and B of AB quartet, *J* = 8.22 Hz, 4H), 7.50 (dd, *J* = 8.41, 2.54 Hz 1H), 7.14 (t, *J* = 8.22 Hz, 1H), 4.56 (s, 2H), 4.17 (m, 3H), 3.91 (s, 3H), 3.70 (t, *J* = 5.67 Hz, 2H), 3.03 (br m, 2H), 3.00 (t, *J* = 5.67 Hz, 2H), 2.90 (s, 3H), 2.40 (br m, 4H), 2.13 (br m, 4H), 1.76 (br d, *J* = 11.15 Hz, 4H).


G. 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid.

To a stirred solution of 635 mg (0.94 mmol) of 6-fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid methyl ester in 10 mL of THF was treated a solution of 53 mg (0.94 mmol) of KOH in 0.5 mL of water. This was stirred overnight, after the hydrolysis was deemed complete 1 mL of 1 *N* HCl solution was added. This was then extracted with EtOAc (3 x 30 mL). The combined organic layers were then washed with brine, dried over Na₂SO₄, and concentrated to yield 622 mg (100%) of a white solid. MS (electrospray): exact mass calculated for C₃₁H₃₂F₄N₄O₄S₂: 664.18; *m/z* found, 665.2 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz): 8.12 (dd, *J* = 8.81, 5.28 Hz, 1H), 7.83 and 7.76 (A and B of AB quartet, *J* = 8.41 Hz, 4H), 7.17 (br t, *J* = 8.61 Hz, 1H), 4.47 (s, 2H), 4.29 (br s, 1H), 4.16 (t, *J* = 7.04 Hz, 2H), 3.53 (t, *J* = 5.67 Hz, 2H), 3.28 (br m, 4H), 3.00 (s, 3H), 2.96 (m, 2H), 2.73 (br s, 2H), 2.52 (br m, 2H), 2.13 (br m, 2H), 1.76 (br m, 2H).

H. 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-hydroxy-ethyl)-amide.

A stirred solution of 20 mg (0.03 mmol) of 6-fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid in 0.3 mL of dry DMF was treated with 14 mg (0.036 mmol) of HBTU and 8 μ L (0.045 mmol) of DIEA. The solution was stirred for 5 min before the addition of 0.01 mL (0.15 mmol) of ethanol amine. The reaction was stirred at room temperature for 30 min then partitioned between EtOAc (30 mL) and saturated NaHCO_3 (20 mL). The aqueous layer was further extracted with EtOAc (2 x 20 mL). The combined organic layers were then washed with brine, dried over Na_2SO_4 , and concentrated. Purification by column chromatography (silica, 5-10% of 2 N NH_3 in $\text{MeOH}/\text{CH}_2\text{Cl}_2$) yielded 16 mg (76%) of a white solid. MS (electrospray): exact mass calculated for $\text{C}_{33}\text{H}_{37}\text{F}_4\text{N}_5\text{O}_4\text{S}_2$: 707.22; m/z found, 708.2 $[\text{M}+\text{H}]^+$. ^1H NMR (CDCl_3 , 400 MHz): 7.97 (br s, 1H), 7.73 and 7.66 (A and B of AB quartet, J = 8.41 Hz, 4H), 7.49 (dd, J = 8.41, 2.35 Hz, 1H), 7.15 (dt, J = 8.61, 2.54 Hz, 1H), 6.41 (t, J = 5.67 Hz, 1H), 4.56 (s, 2H), 4.16 (t, J = 7.04 Hz, 2H), 3.84 (dd, J = 5.28, 4.70 Hz, 2H), 3.70 (t, J = 5.67 Hz, 2H), 3.62 (m, 2H), 3.58 (br s, 1H), 3.05 (br m, 2H), 2.97 (t, J = 5.67 Hz, 2H), 2.91 (s, 3H), 2.40 (br m, 4H), 2.13 (br m, 4H), 1.84 (br d, J = 12.32 Hz, 2H).

20 EXAMPLE 12



6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid (2-amino-ethyl)-amide.

25

A stirred solution of 300 mg (0.43 mmol) 6-fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid in 5 mL of dry DMF was


treated with 812 mg (2.14 mmol) of HBTU and 0.75 mL (4.28 mmol) of DIEA.

The solution was stirred for 5 min before the addition of 0.28 mL (4.28 mmol) of ethylenediamine. The reaction was stirred at room temperature for 30 min then partitioned between EtOAc (30 mL) and saturated sodium bicarbonate (20 mL).

- 5 The aqueous layer was further extracted with EtOAc (2 x 20 mL). The combined organic layers were then washed with brine, dried over Na₂SO₄, and concentrated. Purification by column chromatography (silica, 5-10% of (2 N NH₃ in MeOH)/CH₂Cl₂) afforded 200 mg (66%) of a clear oil. MS (electrospray): exact mass calculated for C₃₃H₃₈F₄N₆O₃S₂: 706.24; *m/z* found, 707.2 [M+H]⁺.
- 10 ¹H NMR (CD₃OD, 400 MHz): 8.14 (m, 1H), 7.84 and 7.72 (A and B of AB quartet, *J* = 8.41 Hz, 4H), 7.68 (m, 1H), 7.21 (dt, *J* = 8.81, 2.74 Hz, 1H), 4.54 (s, 2H), 4.27 (t, *J* = 6.26 Hz, 2H), 4.00-3.90 (m, 2H), 3.76-3.61 (m, 8H), 3.28-3.23 (br m, 1H), 3.19-3.09 (br m, 3H), 2.98 (s, 3H), 2.97-2.93 (m, 2H), 2.67 (br m, 2H), 2.52-2.37 (m, 3H), 2.02 (br d, *J* = 13.89 Hz, 2H).

15

EXAMPLE 13




- 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-morpholin-4-yl-ethyl)-amide.
- 20

- A stirred solution of 20 mg (0.03 mmol) of 6-fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid in 0.3 mL of dry
- 25 DMF was treated with 14 mg (0.036 mmol) of HBTU and 8 μ L (0.045 mmol) of


DIEA. The solution was stirred for 5 min before the addition of 20 μ L (0.15 mmol) of 4-(2-aminoethyl)morpholine. The reaction was stirred at room temperature for 30 min then partitioned between EtOAc (30 mL) and saturated NaHCO_3 (20 mL). The aqueous layer was further extracted with EtOAc (2 x 20 mL). The combined organic layers were then washed with brine, dried over Na_2SO_4 , and concentrated. Purification by column chromatography (silica, 5-10% of (2 N NH_3 in MeOH)/ CH_2Cl_2) afforded 15 mg (65%) of a white solid. MS (electrospray), exact mass calculated for $\text{C}_{37}\text{H}_{44}\text{F}_4\text{N}_6\text{O}_4\text{S}_2$: 776.28; m/z found, 777.3 $[\text{M}+\text{H}]^+$. ^1H NMR (CDCl_3 , 400 MHz): 8.07 (br s, 1H), 7.73 and 7.66 (A and B of AB quartet, $J = 8.41$ Hz, 4H), 7.50 (dd, $J = 8.41$, 2.35 Hz, 1H), 7.14 (dt, $J = 8.61$, 2.54 Hz, 1H), 6.64 (t, $J = 4.70$ Hz, 1H), 4.56 (s, 2H), 4.16 (t, $J = 6.85$ Hz, 2H), 3.80-3.67 (m, 7H), 3.53 (q, $J = 5.48$ Hz, 2H), 3.03 (br m, 2H), 2.98 (t, $J = 5.67$ Hz, 2H), 2.90 (s, 3H), 2.60 (t, $J = 5.87$ Hz, 2H), 2.38 (br m, 4H), 2.12 (br m, 4H), 1.86-1.73 (br m, 3H).

EXAMPLE 14




1-[1-{2-Hydroxy-3-[4-(1H-indol-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone.

EXAMPLE 15




1-[4-(5-Fluoro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-
5 trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-
ol.

EXAMPLE 16




1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-
c]pyridin-1-yl]-3-[4-(5-chloro-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol.

EXAMPLE 17




1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-
20 c]pyridin-1-yl]-3-[4-(5-chloro-2-methyl-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol.

EXAMPLE 18




1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-
pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-methyl-1H-indol-3-yl)-piperidin-1-yl]-propan-
2-ol.

EXAMPLE 19




3-(1-[2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl]-piperidin-4-yl)-1H-indole-5-carbonitrile.

EXAMPLE 20




1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-methoxy-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol.

EXAMPLE 21




3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-
5 tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-
carboxylic acid ethyl ester.

EXAMPLE 22



1-[4-(6-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-
trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-
ol.


EXAMPLE 23



1-[1-(3-{4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-2-hydroxy-propyl)-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone.

5


EXAMPLE 24



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

EXAMPLE 25


15



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

20


EXAMPLE 26



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-
5 pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-
1-yl]-propan-2-ol.

EXAMPLE 27


10



1-[4-(5-Dimethylamino-1H-pyrrolo[3,2-b]pyridin-3-yl)-piperidin-1-yl]-3-[5-
methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-
c]pyridin-1-yl]-propan-2-ol.

15


EXAMPLE 28



1-[4-(5-Dimethylamino-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol.

5


EXAMPLE 29



3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-pyrrolo[2,3-b]pyridine-6-carbonitrile.

EXAMPLE 30


15



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propan-2-ol.

20


EXAMPLE 31



1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-
 5 pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-
 piperidin-1-yl]-propan-2-ol.

EXAMPLE 32


10



1-[4-(6-Fluoro-2-hydroxymethyl-benzo[b]thiophen-3-yl)-piperidin-1-yl]-3-[5-
 methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-
 c]pyridin-1-yl]-propan-2-ol.

15


EXAMPLE 33



6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carbaldehyde.

5


EXAMPLE 34



10 6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid methyl ester.


EXAMPLE 35

15



20 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid amide.

EXAMPLE 36



6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-
5 tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-
2-carboxylic acid ethylamide.

EXAMPLE 37

10

Cathepsin S Inhibition Assay.

Recombinant human cathepsin S (CatS) was expressed in the
baculovirus system and purified in one step with a thiopropyl-sepharose
15 column. 10-L yielded ~700 mg of CatS and N-terminal sequencing confirmed
identity. The assay is run in 100 mM sodium acetate pH 5.0 containing 1 mM
DTT and 100 mM NaCl. The substrate for the assay is

(Aedens)EKARVLAEEA(Dabcyl)K-amide

20

The K_m for the substrate is around 5 μ M but the presence of substrate inhibition
makes kinetic analysis difficult. With 20 μ M substrate the assay rate is linear
over the range of 1-8 ng CatS in 100 μ l reaction. Using 2 ng/well of CatS, the
production of product is linear and yields ~7-fold signal after 20 min with only
25 20% loss of substrate. Primary assays are run by quenching the reaction after
20 min with 0.1% SDS and then measuring the fluorescence. For other
assays, measurements are taken every min for 20 min. The rate is calculated
from the slope of the increase and the percent inhibition is calculated from this
(See Tables 1, 2 and 3 below).

Table 1

EXAMPLE	IC ₅₀ (μM)
1	0.07
2	0.03
3	0.05
4	0.09
5	0.03
6	0.03
7	0.05
8	0.03
9	0.02
10	0.02
11	0.02
12	0.05
13	0.05

Table 2

EXAMPLE	IC ₅₀ (μM)
14	0.07
15	0.04
16	0.06
17	0.03
18	0.06
19	0.02
20	0.02
21	0.04
22	0.03
23	0.08
24	0.13
25	0.05
26	0.09
27	0.10
28	0.07
29	0.08
30	0.02
31	0.07
32	0.14
33	0.08
34	0.13
35	0.03
36	0.04

EXAMPLE 101

1-(3-(4-Chloro-3-methyl-phenyl)-1-{2-hydroxy-3-[4-(1H-indol-3-yl)-piperidin-1-yl]-propyl}-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl)-ethanone

5 EXAMPLE 102

1-[1-{3-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-2-hydroxy-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

EXAMPLE 103

10 1-{3-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-propyl}-5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine

EXAMPLE 104

15 3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-carbonitrile

EXAMPLE 105

20 1-[4-(6-Chloro-1-methyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol

EXAMPLE 106

25 1-[4-(6-Chloro-1-methanesulfonyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol

EXAMPLE 107

30 1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-chloro-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol

EXAMPLE 108

3-(1-{3-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-1H-indole-5-carbonitrile

5 EXAMPLE 109

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-{4-[6-chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-propan-2-ol

10 EXAMPLE 110

1-[3-(4-Bromo-phenyl)-1-(3-{4-[6-chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-2-hydroxy-propyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

15 EXAMPLE 111

1-[1-{2-Hydroxy-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

EXAMPLE 112

20 1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol

EXAMPLE 113

25 1-(3-(4-Bromo-phenyl)-1-{2-hydroxy-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propyl}-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl)-ethanone

EXAMPLE 114

30 1-[1-(2-Hydroxy-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propyl)-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

EXAMPLE 115

5-Methanesulfonyl-1-{3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-3,6-dihydro-2H-pyridin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine

5

EXAMPLE 116

5-Methanesulfonyl-1-{3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine

10 EXAMPLE 117

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propan-2-ol

15 EXAMPLE 118

1-[3-(4-Bromo-phenyl)-1-(2-hydroxy-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

20 EXAMPLE 119

6-Chloro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-pyrrolo[2,3-b]pyridine-1-carboxylic acid methyl ester

25 EXAMPLE 120

1-[4-(6-Chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol

30 EXAMPLE 121

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-oxy-1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol

EXAMPLE 122

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol

5

EXAMPLE 123

1-[1-{2-Hydroxy-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

10 EXAMPLE 124

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol

EXAMPLE 125

15 1-(3-(4-Bromo-phenyl)-1-{2-hydroxy-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propyl}-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl)-ethanone

EXAMPLE 126

20 3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-5-oxy-pyrrolo[3,2-c]pyridine-1-carboxylic acid methyl ester

EXAMPLE 128

25 3-(1-{3-[5-Acetyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-6-fluoro-benzofuran-2-carboxylic acid methyl ester

EXAMPLE 129

30 3-(1-{3-[5-Acetyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-6-fluoro-benzo[b]thiophene-2-carboxylic acid methyl ester

EXAMPLE 130

1-[4-(6-Fluoro-benzo[b]thiophen-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol

5

EXAMPLE 131

1-[1-{3-[4-(6-Fluoro-benzofuran-3-yl)-piperidin-1-yl]-2-hydroxy-propyl}-3-(4trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone

10

EXAMPLE 132

6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid

15

EXAMPLE 133

6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid dimethylamide

20

EXAMPLE 134

3-(1-{3-[5-Acetyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-6-fluoro-benzo[b]thiophene-2-carbonitrile

25

EXAMPLE 135

6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carbonitrile

30

Table 3

EXAMPLE	IC ₅₀ (μM)
101	0.25
102	0.11
103	0.16
104	0.14
105	0.63
106	0.20
107	0.04
108	0.04
109	0.08
110	0.14
111	0.17
112	0.13
113	0.20
114	0.11
115	0.59
116	0.25
117	0.10
118	0.26
119	0.69
120	0.25
121	0.31
122	0.19
123	0.15
124	0.13
125	0.21
126	0.25
127	0.40
128	0.23

EXAMPLE (Table 3, cont'd)	IC ₅₀ (μM) (Table 3, cont'd)
130	0.19
131	0.37
132	0.09
133	0.38
134	0.38
135	0.21

F. Other Embodiments

The features and advantages of the invention are apparent to one of ordinary skill in the art. Based on this disclosure, including the summary, detailed description, background, examples, and claims, one of ordinary skill in the art will be able to make modifications and adaptations to various conditions and usages. These other embodiments are also within the scope of the invention.


10

What is claimed is:

CLAIMS

1. A compound of formula (I):

5



(I)

wherein:

- 10 the dashed line adjacent C-R⁶ is absent or an sp² bond;
- Y is nitrogen or R²⁰C;
- Z is nitrogen or R²¹C;
- T is nitrogen or R²C;
- S is nitrogen or R³C;
- 15 provided between 0 and 3 of S, T, Y, and Z are nitrogen; and further provided that one of S, T, Y, and Z can be =N⁺-O⁻ where the remaining three are not nitrogen;
- R²⁰ is selected from hydrogen, halogen, C₁₋₅ alkoxy, hydroxy, C₁₋₅ alkyl, cyano, nitro, C₁₋₅ haloalkyl, R^cR^pN, R^cR^pNC=O, C₂₋₈ acyl, 4-7 membered heterocyclyl, (4-7 membered heterocyclyl)-C₁₋₅ alkylene, phenyl, (phenyl)C₁₋₅ alkylene, R¹⁴OC=O, R¹⁴S, R¹⁴SO, and R¹⁴SO₂;
- 20 R²¹ is selected from hydrogen, halogen, C₁₋₅ alkoxy, hydroxy, C₁₋₅ alkyl, cyano, nitro, C₁₋₅ haloalkyl, R^cR^dN, R^cR^dNC=O, C₂₋₈ acyl, 4-7 membered heterocyclyl, (4-7 membered heterocyclyl)-C₁₋₅ alkylene, phenyl, (phenyl)C₁₋₅ alkylene, R¹⁵OC=O, R¹⁵S, R¹⁵SO, and R¹⁵SO₂;
- 25

- R^2 is selected from hydrogen, halogen, C_{1-5} alkoxy, hydroxy, C_{1-5} alkyl, cyano, nitro, C_{1-5} haloalkyl, $R^e R^f N$, $R^e R^f NC=O$, C_{2-8} acyl, 4-7 membered heterocyclyl, (4-7 membered heterocyclyl)- C_{1-5} alkylene, phenyl, (phenyl) C_{1-5} alkylene, $R^{16}OC=O$, $R^{16}S$, $R^{16}SO$, and $R^{16}SO_2$;
- 5 R^3 is selected from hydrogen, halogen, C_{1-5} alkoxy, hydroxy, C_{1-5} alkyl, cyano, nitro, C_{1-5} haloalkyl, $R^g R^h N$, C_{2-8} acyl, 4-7 membered heterocyclyl, (4-7 membered heterocyclyl)- C_{1-5} alkylene, phenyl, (phenyl) C_{1-5} alkylene, $R^{17}OC=O$, $R^m R^n NC=O$, $R^m R^n NSO_2$, $R^{17}S$, $R^{17}SO$, and $R^{17}SO_2$;
- R^5 and R^6 are independently selected from hydrogen and C_{1-5} alkyl;
- 10 R^7 and R^8 independently are hydrogen, C_{1-5} alkyl, C_{1-5} alkenyl, C_{1-5} alkoxy, C_{1-5} alkylthio, halogen, or 4-7 membered carbocyclyl or heterocyclyl; alternatively, R^7 and R^8 can be taken together to form an optionally substituted 5- to 7- membered carbocyclic or heterocyclic ring, which ring may be unsaturated or aromatic; said ring being optionally
- 15 substituted with between 1 and 3 substituents independently selected from halo, hydroxy, cyano, nitro, amino, R^i , R^iO- , R^iS- , $R^iO(C_{1-5} alkylene)-$, $R^iO(C=O)-$, $R^i(C=O)-$, $R^i(C=S)-$, $R^i(C=O)O-$, $R^iO(C=O)(C=O)-$, R^iSO_2 , $NHR^u(C=NH)-$, NHR^uSO_2- , and $NHR^u(C=O)-$;
- R^i is C_{1-6} alkyl, phenyl, benzyl, phenethyl, or C_{2-5} heterocyclyl, (C_{1-5} heterocyclyl) C_{1-6} alkylene, NH_2 , mono- or di(C_{1-6} alkyl) $N-$, or $R^{49}OR^{50}-$,
 20 wherein R^{49} is H, C_{1-5} alkyl, C_{2-5} alkenyl, phenyl, benzyl, phenethyl, C_{1-5} heterocyclyl, or (C_{1-5} heterocyclyl) C_{1-6} alkylene and R^{50} is C_{1-5} alkylene, phenylene, or divalent C_{1-5} heterocyclyl; and
- R^u can be H in addition to the values for R^i ;
- 25 R^c is hydrogen, C_{1-5} alkyl, phenyl, C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl, $R^{10}OC=O-$, $R^i R^j NC=O$, $R^{10}SO-$, $R^{10}SO_2-$, and $R^i R^j NSO_2$;
- R^e is hydrogen, C_{1-5} alkyl, phenyl, C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl, $R^{40}OC=O$, $R^{43}R^{44}NC=O$, $R^{40}SO$, $R^{40}SO_2$, and $R^{43}R^{44}NSO_2$;
- R^m is hydrogen, C_{1-5} alkyl, phenyl, C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl,
 30 $R^{41}OC=O$, $R^{45}R^{46}NC=O$, $R^{41}SO$, $R^{41}SO_2$, and $R^{45}R^{46}NSO_2$;
- R^o is hydrogen, C_{1-5} alkyl, phenyl, C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl, $R^{42}OC=O$, $R^{47}R^{48}NC=O$, $R^{42}SO$, $R^{42}SO_2$, and $R^{47}R^{48}NSO_2$;

each of R^d , R^f , R^n , and R^p is independently selected from hydrogen, C_{1-5} alkyl, phenyl, and C_{2-5} heterocyclyl; in addition, R^c and R^d , R^e and R^f , R^m and R^n , or R^o and R^p , independently, can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring
5 may be saturated, unsaturated or aromatic;

each of R^9 , R^{10} , R^{11} , R^{14} , R^{15} , R^{16} , R^{17} , R^{40} , R^{41} , and R^{42} is independently C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl;

each of R^i and R^j , R^k and R^l , R^{43} and R^{44} , R^{45} and R^{46} , R^{47} and R^{48} are independently hydrogen, C_{1-5} alkyl, C_{3-5} alkenyl, phenyl, or C_{2-5} heterocyclyl; in addition, R^i and R^j , and R^k and R^l , R^{43} and R^{44} , R^{45} and R^{46} , and R^{47} and R^{48} , independently, can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring
10 may be saturated, unsaturated or aromatic;

R^9 is hydrogen, C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl, C_{2-8} acyl, aroyl, $R^9OC=O$, $R^{18}R^{19}NC=O$, R^9SO , R^9SO_2 , or $R^{18}R^{19}NSO_2$;

R^h is hydrogen, C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl; alternatively, R^g and R^h can be taken together to form an optionally substituted 4- to 7- membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

R^{18} and R^{19} independently are hydrogen, C_{1-5} alkyl, phenyl, or C_{2-5} heterocyclyl; alternatively, R^{18} and R^{19} can be taken together to form an optionally substituted 4- to 7- membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

n is 0, 1 or 2;

G is C_{3-6} alkenediyl or C_{3-6} alkanediyl, optionally substituted with hydroxy, halogen, C_{1-5} alkyl, C_{1-5} alkoxy, oxo, hydroximino, CO_2R^k , NR^kR^l , (L)- C_{1-4} alkylene-, $R^kR^lNCO_2$, [(L)- C_{1-5} alkylene]amino, N_3 , or (L)- C_{1-5} alkoxy;

L is amino, mono- or di- C_{1-5} alkylamino, pyrrolidinyl, morpholinyl, piperidinyl, homopiperidinyl, or piperazinyl, wherein available ring nitrogens can be optionally substituted with C_{1-5} alkyl, benzyl, C_{2-5} acyl, C_{1-5} alkylsulfonyl, or C_{1-5} alkoxycarbonyl;

Ar represents a monocyclic or bicyclic aryl or heteroaryl ring, optionally substituted with between 1 and 3 substituents independently selected

from halogen, C₁₋₅ alkoxy, C₁₋₅ alkyl, C₂₋₅ alkenyl, cyano, azido, nitro, R²²R²³N, R²²S, R²²SO, R²²SO₂, R²²OC=O, R²²R²³NC=O, C₁₋₅ haloalkyl, C₁₋₅ haloalkoxy, C₁₋₅ haloalkylthio, and C₁₋₅ alkylthio;

5 R²² is hydrogen, C₁₋₅ alkyl, C₃₋₅ alkenyl, phenyl, benzyl, C₂₋₅ heterocyclyl, C₂₋₈ acyl, aroyl, R¹¹OC=O, R²⁴R²⁵NC=O, R¹¹S, R¹¹SO, R¹¹SO₂, or R²⁴R²⁵NSO₂;

R²³ is hydrogen, C₁₋₅ alkyl, phenyl, benzyl, or C₂₋₅ heterocyclyl; alternatively, R²² and R²³ can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

10 R²⁴ and R²⁵ are independently hydrogen, C₁₋₅ alkyl, phenyl, benzyl, or C₁₋₅ heteroaryl;

alternatively, R²⁴ and R²⁵ can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

15 R³² is hydrogen, C₁₋₅ alkyl, cyano, C₁₋₅ hydroxyalkyl, C₂₋₈ acyl, -(C=O)NR^vR^x, CHO, or C₁₋₆ alkoxycarbonyl, wherein each of R^v and R^x is independently selected from H, C₁₋₅ alkyl, C₁₋₅ hydroxyalkyl, C₁₋₅ heterocyclyl, (C₁₋₅ heterocyclyl) C₁₋₅ alkylene, C₁₋₅ aminoalkylene, C₃₋₈ acyloxy, CHO, C₁₋₆ alkoxycarbonyl, and cyano;

20 Q is NR³³, S, or O;

R³³ represents hydrogen, C₁₋₅ alkyl, phenyl, benzyl, phenethyl, C₂₋₅ heterocyclyl, (C₂₋₅ heterocyclyl)C₁₋₅ alkylene, C₂₋₈ acyl, aroyl, R³⁵OC=O, R³⁶R³⁷NC=O, R³⁵SO, R³⁵S, R³⁵SO₂ and R³⁶R³⁷NSO₂;

25 R³⁵ is selected from hydrogen, C₁₋₅ alkyl, phenyl, benzyl, phenethyl, and C₂₋₅ heteroaryl;

R³⁶ and R³⁷ are each independently selected from hydrogen, C₁₋₅ alkyl, phenyl, or C₂₋₅ heteroaryl;

30 alternatively, R³⁶ and R³⁷ can be taken together to form an optionally substituted 4- to 7-membered ring heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

wherein each of the above hydrocarbyl or heterocarbyl groups, unless otherwise indicated, and in addition to any specified substituents, is optionally and independently substituted with between 1 and 3 substituents selected from methyl, halomethyl, hydroxymethyl, halo, hydroxy, amino, nitro, cyano, C₁₋₅ alkyl, C₁₋₅ alkoxy, -COOH, C₂₋₆ acyl, [di(C₁₋₄ alkyl)amino]C₂₋₅ alkylene, [di(C₁₋₄ alkyl)amino] C₂₋₅ alkyl-NH-CO-, and C₁₋₅ haloalkoxy;

or a pharmaceutically acceptable salt, amide, or ester thereof; or a stereoisomeric form thereof.

2. A compound of claim 1, wherein one of S, T, Y, and Z is nitrogen.
3. A compound of claim 1, wherein S and T are CR³ and CR², respectively.
4. A compound of claim 1, wherein S, T, Y, and Z are CR³, CR², CR²⁰, and CR²¹, respectively.
5. A compound of claim 1, wherein:
 - (a) Z is N, Y is N, S is CR³, and T is CR²; or
 - (b) S is N, T is N, Y is CR²⁰, and Z is CR²¹.
6. A compound of claim 1, wherein R² is hydrogen, halogen, C₁₋₅ alkoxy, cyano, R^eR^fN, or a 5-6 membered heterocyclyl.
7. A compound of claim 1, wherein R³ is hydrogen, halogen, C₁₋₅ alkoxy, C₁₋₅ alkyl, cyano, R¹⁷OC=O, or R^gR^hN, where R^g and R^h are H or C₁₋₅ alkyl, or are taken together to form a 5-6 membered heterocyclyl.
8. A compound of claim 1, wherein each of R² and R³ is independently selected from hydrogen, halogen, and a 5-6 membered heterocyclyl.

9. A compound of claim 1, wherein R^5 and R^6 are independently selected from hydrogen and C_{1-3} alkyl.
10. A compound of claim 9, wherein one of R^5 and R^6 is H.
11. A compound of claim 10, wherein R^5 and R^6 are each H.
12. A compound of claim 1, wherein one of R^7 and R^8 is H and the other is 5-7 membered carbocyclyl or heterocyclyl.
13. A compound of claim 1, wherein R^7 and R^8 are taken together to form an optionally substituted 5- to 7- membered carbocyclic or heterocyclic ring.
14. A compound of claim 13, wherein R^7 and R^8 are taken together to form a six-membered heterocyclyl.
15. A compound of claim 13 wherein R^7 and R^8 taken together form a 5-7 membered heterocyclyl optionally N-substituted with $R^t(C=O)-$, R^tSO_2- , or $NHR^u(C=O)-$ wherein R^t is C_{1-6} alkyl, phenyl, or C_{2-5} heterocyclyl and R^u is H, C_{1-6} alkyl, phenyl, or C_{2-5} heterocyclyl.
16. A compound of claim 1, wherein each of R^c , R^e , R^m , and R^o is independently selected from hydrogen, C_{1-5} alkyl, C_{2-8} acyl, $(C_{1-5} \text{ alkyl})OC=O$, and the respective $RRNC=O$, RSO , RSO_2 , and $RRNSO_2$ groups.
17. A compound of claim 1, wherein each of R^c , R^d , R^g , R^h , R^o , R^f , and R^p is independently selected from hydrogen and C_{1-5} alkyl; or, independently, R^e and R^f , R^g and R^h , or R^o and R^p taken together form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring.
18. A compound of claim 17 wherein R^e and R^f taken together are morpholinyl, piperidinyl, or pyrrolidinyl.

19. A compound of claim 1, wherein each of R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^i , R^j , R^k and R^l independently is hydrogen or C_{1-5} alkyl.
- 5 20. A compound of claim 1, wherein each of R^9 , R^{11} , R^{14} , R^{15} , R^{16} and R^{17} is independently C_{1-5} alkyl.
- 10 21. A compound of claim 1, wherein R^9 is C_{1-5} alkyl, C_{2-8} acyl, $R^9OC=O$, $R^{18}R^{19}NC=O$, R^9SO , R^9SO_2 , or $R^{18}R^{19}NSO_2$; and R^h is H or C_{1-5} alkyl; alternatively, R^9 and R^h can be taken together to form an optionally substituted 5- to 6-membered heterocyclyl.
22. A compound of claim 21, wherein R^9 and R^h are each C_{1-3} alkyl.
- 15 23. A compound of claim 1, wherein R^{18} and R^{19} independently are hydrogen or C_{1-5} alkyl.
24. A compound of claim 1, wherein n is 1.
- 20 25. A compound of claim 1, wherein G is C_{3-4} alkanediyl, optionally substituted with hydroxy, halogen, $[(L)-C_{1-5}$ alkylene]amino, or $(L)-C_{1-5}$ alkyloxy.
- 25 26. A compound of claim 25, wherein G is C_3 alkanediyl, optionally substituted with hydroxy.
- 30 27. A compound of claim 1, wherein R^{20} and R^{21} are independently selected from hydrogen, halogen, C_{1-5} alkoxy, C_{1-5} alkyl, cyano, nitro, 4-7 membered heterocyclyl, and R^oR^pN or R^cR^dN , respectively.
28. A compound of claim 27, wherein R^{20} and R^{21} are independently selected from hydrogen, halogen, 5- to 6-membered heterocyclyl, and R^oR^pN or R^cR^dN , respectively.

29. A compound of claim 1, wherein Ar represents a monocyclic ring, optionally substituted with 1 to 2 substituents selected from halogen, C₁₋₅ alkyl, cyano, nitro, R²²R²³N, C₁₋₃ haloalkyl, and C₁₋₃ haloalkoxy.
30. A compound of claim 29, wherein Ar is a six-membered aromatic ring monosubstituted at the 4-position with halogen, methyl, CF₃, or OCF₃, or disubstituted at the 3- and 4-positions with substituents independently selected from halogen, CF₃, methyl, and OCF₃.
31. A compound of claim 29, wherein each of R²², R²³, and R²⁴ is independently hydrogen or C₁₋₅ alkyl.
32. A compound of claim 1, wherein R²⁵ and R²⁶ independently are hydrogen or C₁₋₅ alkyl, or, alternatively, R²⁵ and R²⁶ are taken together to form an optionally substituted 4- to 7- membered heterocyclic ring, which ring may be saturated, unsaturated or aromatic.
33. A compound of claim 32, wherein R²⁵ and R²⁶ independently are hydrogen or C₁₋₅ alkyl.
34. A compound of claim 1, wherein Q is NR³³ or S.
35. A compound of claim 34, wherein Q is NR³³, R³³ is H or C₂₋₅ heterocyclyl, and R³² is H, C₁₋₅ alkyl, C₁₋₅ hydroxyalkyl, -(C=O)NR^vR^x, CHO, or C₁₋₆ alkoxy carbonyl, wherein each of R^v and R^x is independently selected from H, C₁₋₅ hydroxyalkyl, (C₁₋₅ heterocyclyl)-C₁₋₅ alkylene, and C₁₋₅ aminoalkylene.
36. A compound of claim 34, wherein Q is S and R³³ is NR³⁶R³⁷(C=O)- where each of R³⁶ and R³⁷ are independently selected from hydrogen and C₁₋₅ alkyl.

37. A compound of claim 1, wherein R^{35} is selected from hydrogen and C_{1-5} alkyl; R^{36} and R^{37} are each independently selected from hydrogen, C_{1-5} alkyl, or, alternatively, R^{36} and R^{37} can be taken together to form an optionally substituted 4- to 7-membered heterocyclic ring.
38. A compound of claim 1, wherein
- Y is nitrogen or $R^{20}C$;
- Z is nitrogen or $R^{21}C$;
- 10 T is nitrogen or $R^{22}C$;
- S is nitrogen or $R^{23}C$;
- provided between 0 and 2 of S, T, Y, and Z are nitrogen;
- R^2 is hydrogen, halogen, hydroxy, C_{1-5} alkoxy, C_{1-5} alkyl, 5- to 6-membered heterocyclyl, or R^eR^fN ;
- 15 R^3 is hydrogen, halogen, C_{1-5} alkoxy, hydroxy, C_{1-5} alkyl, , 5- to 6-membered heterocyclyl, or R^gR^hN ;
- R^5 and R^6 are each H;
- R^7 and R^8 independently are taken together to form an optionally substituted 5- to 7- membered unsaturated heterocyclic ring;
- 20 each of R^a , R^e , R^m , and R^o is independently selected from hydrogen, C_{1-5} alkyl, C_{2-8} acyl, $(C_{1-5} \text{ alkyl})OC=O$, and the respective $RRNC=O$, RSO , RSO_2 , and $RRNSO_2$ groups;
- each of R^b , R^f , R^n , and R^p , is independently selected from hydrogen and C_{1-5} alkyl;
- 25 each of R^9 , R^{11} , R^{14} , R^{15} , R^{16} , R^{17} , R^{40} , R^{41} and R^{42} is independently C_{1-5} alkyl;
- each of R^c , R^d , R^i , R^j , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^k and R^l are independently are hydrogen or C_{1-5} alkyl;
- R^9 is hydrogen, or C_{1-5} alkyl, C_{2-8} acyl, $R^9OC=O$, $R^{18}R^{19}NC=O$, R^9SO , R^9SO_2 , or $R^{18}R^{19}NSO_2$;
- 30 R^h is hydrogen or C_{1-5} alkyl;
- alternatively, R^9 and R^h can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

R^{18} and R^{19} independently are hydrogen or C_{1-5} alkyl;

n is 0 or 1;

G is C_{3-4} alkenediyl or C_{3-4} alkanediyl, optionally substituted with hydroxy, halogen, C_{1-5} alkyloxy, oxo, hydroximino, CO_2R^k , $R^kR^lNCO_2$, N_3 , or $(L)-C_{1-5}$ alkoxy;

L is amino, mono- or di- C_{1-5} alkylamino, pyrrolidinyl, morpholinyl, piperidinyl, homopiperidinyl, or piperazinyl, available ring nitrogens being optionally with C_{1-5} alkyl, benzyl, C_{2-5} acyl, or C_{1-5} alkyloxycarbonyl;

R^{20} and R^{21} are independently selected from hydrogen, halogen, C_{1-5} alkoxy,

C_{1-5} alkyl, cyano, nitro, and R^oR^pN ;

alternatively, R^3 and R^{20} or R^3 and R^{21} can be taken together to form an optionally substituted 5- or 6-membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

Ar represents a monocyclic or bicyclic aryl or heteroaryl ring, optionally

substituted with hydrogen, halogen, C_{1-5} alkoxy, C_{1-5} alkyl, cyano, nitro, $R^{22}R^{23}N$, $R^{24}SO_2$, $R^{24}OC=O$, $R^{25}R^{26}NC=O$, CF_3 , OCF_3 , SCF_3 , or C_{1-5} alkylthio;

R^{22} is hydrogen, C_{1-5} alkyl, phenyl, benzyl, phenethyl, C_{2-5} heteroaryl, C_{2-8} acyl, aroyl, $R^{24}OC=O$, $R^{25}R^{26}NC=O$, $R^{24}SO$, $R^{24}SO_2$, or $R^{25}R^{26}NSO_2$;

R^{23} is hydrogen or C_{1-5} alkyl;

alternatively, R^{22} and R^{23} can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

R^{24} is hydrogen or C_{1-5} alkyl;

R^{25} and R^{26} are independently hydrogen or C_{1-5} alkyl;

or, alternatively, R^{25} and R^{26} can be taken together to form an optionally substituted 4- to 7- membered carbocyclic or heterocyclic ring, which ring may be saturated, unsaturated or aromatic;

R^{32} is hydrogen, C_{1-5} alkyl, C_{1-5} hydroxyalkyl, CHO , C_{2-6} acyl, C_{1-6}

alkoxycarbonyl, or $-(C=O)NR^vR^x$, wherein each of R^vR^x is independently selected from H , C_{1-5} alkyl, C_{1-5} hydroxyalkyl, C_{3-8} acyloxy, (amino) C_{1-6} alkylene, $(C_{1-5} \text{ heterocycl})C_{1-5}$ alkylene, or C_{1-6} alkoxycarbonyl;

Q is NR^{33} or S ;

R^{33} represents hydrogen, C_{1-5} alkyl, phenyl, benzyl, (C_{2-5} heterocyclyl) C_{1-5} alkylene, C_{2-8} acyl, aroyl, $R^{35}OC=O$, $R^{36}R^{37}NC=O$, $R^{35}SO_2$ and $R^{36}R^{37}NSO_2$;

R^{35} is selected from hydrogen and C_{1-5} alkyl;

5 R^{36} and R^{37} are each independently selected from hydrogen and C_{1-5} alkyl.

39. A compound of claim 1, wherein

one of R^5 and R^6 is H,

10 R^7 and R^8 are taken together to form an optionally substituted 6-membered carbocyclic or heterocyclic ring; and

Ar represents a monocyclic ring, optionally substituted with 1 to 2 substituents selected from halogen, C_{1-5} alkyl, cyano, nitro, $R^{22}R^{23}N$, CF_3 and OCF_3 .

15 40. A compound of claim 39, wherein

both R^5 and R^6 are each H, and

Ar is a six membered ring substituted with halogen, CF_3 , methyl, halomethyl, or OCF_3 , at the 3- or 4- position, or disubstituted at the 3- and 4- positions.

20

41. A compound of claim 40, wherein R^7 and R^8 taken together form pyridinyl, pyrimidinyl, or piperazinyl, optionally N-substituted with $-(C=O)R^t$, SO_2-R^t , or $-(C=O)NHR^u$.

25 42. A compound of claim 39, wherein R^{22} and R^{23} taken together are independently morpholinyl, piperidyl, or pyrrolidinyl, optionally substituted.

30 43. A compound of claim 1, wherein the dashed line adjacent C- R^6 is absent.

44. A compound of claim 1, selected from:

1-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[4-(7-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

10 1-[4-(5-Chloro-2-methyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

20 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

25 1-[4-(6-Dimethylamino-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

30 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-hydroxy-ethyl)-amide;

5 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-amino-ethyl)-amide; and

6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-morpholin-4-yl-ethyl)-amide.

10

45. A compound of claim 1, selected from:

1-[1-{2-Hydroxy-3-[4-(1H-indol-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

15 1-[4-(5-Fluoro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ;

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-chloro-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ;

20 1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-chloro-2-methyl-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ;

25 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-methyl-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ;

3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-carbonitrile ;

30 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-methoxy-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol ;

3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-carboxylic acid ethyl ester ;

1-[4-(6-Chloro-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ;

1-[1-(3-{4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-2-hydroxy-propyl)-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone ;

10 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ;

15 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ;

20 1-[4-(5-Dimethylamino-1H-pyrrolo[3,2-b]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ;

1-[4-(5-Dimethylamino-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ;

25 3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-pyrrolo[2,3-b]pyridine-6-carbonitrile ;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propan-2-ol ;

30 1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-morpholin-4-yl-1H-pyrrolo[2,3-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol ;

1-[4-(6-Fluoro-2-hydroxymethyl-benzo[b]thiophen-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol ;

5 6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carbaldehyde ;

6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid methyl ester ;

10 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid amide ;

and 6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid ethylamide.

46. A compound of claim 1, selected from:

1-{4-[6-Chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)benzo[b]thiophene-2-carboxylic acid (2-morpholin-4-yl-ethyl)-amide;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(5-oxy-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol; and

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol.

47. A compound of claim 1, selected from:

1-(3-(4-Chloro-3-methyl-phenyl)-1-{2-hydroxy-3-[4-(1H-indol-3-yl)-piperidin-1-yl]-propyl}-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl)-ethanone;

1-[1-{3-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-2-hydroxy-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

1-{3-[4-(5-Chloro-1H-indol-3-yl)-piperidin-1-yl]-propyl}-5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine;

3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-1H-indole-5-carbonitrile;

1-[4-(6-Chloro-1-methyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[4-(6-Chloro-1-methanesulfonyl-1H-indol-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-chloro-1H-indol-3-yl)-piperidin-1-yl]-propan-2-ol;

3-(1-{3-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-1H-indole-5-carbonitrile;

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-{4-[6-chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-propan-2-ol;

1-[3-(4-Bromo-phenyl)-1-(3-{4-[6-chloro-1-(2-morpholin-4-yl-ethyl)-1H-indol-3-yl]-piperidin-1-yl}-2-hydroxy-propyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

1-[1-{2-Hydroxy-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

1-(3-(4-Bromo-phenyl)-1-{2-hydroxy-3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propyl}-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl)-ethanone;

1-[1-(2-Hydroxy-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propyl)-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-
5 pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

5-Methanesulfonyl-1-{3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-3,6-dihydro-2H-pyridin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine; and

5-Methanesulfonyl-1-{3-[4-(1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[4,3-
10 c]pyridine.

48. A compound of claim 1, selected from:

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-
15 pyrazolo[4,3-c]pyridin-1-yl]-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propan-2-ol;

1-[3-(4-Bromo-phenyl)-1-(2-hydroxy-3-{4-[1-(2-morpholin-4-yl-ethyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-piperidin-1-yl}-propyl)-1,4,6,7-tetrahydro-
pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

20 6-Chloro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-pyrrolo[2,3-b]pyridine-1-carboxylic acid methyl ester;

1-[4-(6-Chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-
25 c]pyridin-1-yl]-propan-2-ol;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(7-oxy-1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

1-[5-Methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-
30 pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(6-morpholin-4-yl-1H-pyrrolo[2,3-b]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

1-[1-{2-Hydroxy-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

1-[3-(4-Bromo-phenyl)-5-methanesulfonyl-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propan-2-ol;

1-(3-(4-Bromo-phenyl)-1-{2-hydroxy-3-[4-(1H-pyrrolo[3,2-c]pyridin-3-yl)-piperidin-1-yl]-propyl}-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl)-ethanone;

3-(1-{2-Hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-5-oxy-pyrrolo[3,2-c]pyridine-1-carboxylic acid methyl ester;

3-(1-{3-[5-Acetyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-6-fluorobenzofuran-2-carboxylic acid methyl ester;

3-(1-{3-[5-Acetyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-6-fluorobenzo[b]thiophene-2-carboxylic acid methyl ester;

1-[4-(6-Fluoro-benzo[b]thiophen-3-yl)-piperidin-1-yl]-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propan-2-ol;

1-[1-{3-[4-(6-Fluoro-benzofuran-3-yl)-piperidin-1-yl]-2-hydroxy-propyl}-3-(4-trifluoromethyl-phenyl)-1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-5-yl]-ethanone;

6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid;

6-Fluoro-3-(1-{3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carboxylic acid dimethylamide;

3-(1-{3-[5-Acetyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-2-hydroxy-propyl}-piperidin-4-yl)-6-fluorobenzo[b]thiophene-2-carbonitrile; and

6-Fluoro-3-(1-{2-hydroxy-3-[5-methanesulfonyl-3-(4-trifluoromethyl-phenyl)-4,5,6,7-tetrahydro-pyrazolo[4,3-c]pyridin-1-yl]-propyl}-piperidin-4-yl)-benzo[b]thiophene-2-carbonitrile.

- 5 49. A pharmaceutical composition, comprising a compound of claim 1, 38, 44, 45, or 46 and a pharmaceutically acceptable carrier.
- 10 50. A method for treating a subject with a condition mediated by cathepsin S, said method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of claim 1, 38, 44, 45, or 46.
- 15 51. A method for inhibiting cathepsin S activity in a subject, said method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of claim 1, 27, 44, or 46.
- 20 52. A method for treating an autoimmune disease, or inhibiting the progression of an autoimmune disease, in a subject, said method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of claim 1, 27, 44 or 46.
- 25 53. A method of claim 52, wherein the autoimmune disease is selected from lupus, rheumatoid arthritis, and asthma.
- 30 54. A method of claim 52, wherein the autoimmune disease is asthma.
55. A method for treating or inhibiting the progression of tissue transplant rejection in a subject, said method comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising a compound of claim 1, 27, 44 or 46.

56. A method of claim 55, wherein said administration occurs after said subject has undergone a tissue transplant procedure.
57. A method of claim 55, wherein said administration to said subject occurs before or during a tissue transplant procedure.

10

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 February 2002 (21.02.2002)

PCT

(10) International Publication Number
WO 02/014317 A3

(51) International Patent Classification⁷: C07D 471/04,
519/00, A61K 31/437, A61P 29/00, 11/06 // (C07D
471/04, 231:00, 221:00) (C07D 519/00, 471:00, 471:00)

(74) Agents: JOHNSON, Philip, S. et al.; One Johnson &
Johnson Plaza, New Brunswick, NJ 08933 (US).

(21) International Application Number: PCT/US01/25180

(22) International Filing Date: 10 August 2001 (10.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/225,178 14 August 2000 (14.08.2000) US
09/927,188 10 August 2001 (10.08.2001) US

(71) Applicant: ORTHO MCNEIL PHARMACEUTICAL,
INC. [US/US]; U.S. Route 202, P.O. Box 300, Raritan, NJ
08869-0602 (US).

(72) Inventors: CAI, Hui; 11213 Corte Belleza, San Diego,
CA 92130 (US). EDWARDS, James, P.; 8723 Hesby
Court, San Diego, CA 92129 (US). MEDUNA, Steven,
P.; 9975 Paseo Montrail, San Diego, CA 92129 (US). PIO,
Barbara, A.; 3898 Ashford Street, San Diego, CA 92111
(US). WEI, Jianmei; 12516 Senda Panacea, San Diego,
CA 92129 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).


Published:

— with international search report

(88) Date of publication of the international search report:
4 July 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED PYRAZOLES



(57) Abstract: Substituted pyrazoles, methods of manufacturing them, compositions containing them, and methods of using them to treat, for example, autoimmune diseases mediated by cathepsin S are described.

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/US 01/25180

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D471/04 C07D519/00 A61K31/437 A61P29/00 A61P11/06 //(C07D471/04,231:00,221:00),(C07D519/00,471:00,471:00)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) CHEM ABS Data, EPO-Internal, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; NAKATSUKA, MASASHI ET AL: "Preparation of pyrazole derivatives as immunosuppressants" retrieved from STN Database accession no. 130:52417 XP002193692 abstract -& WO 98 56785 A (SUMITOMO PHARMACEUTICALS CO., LTD., JAPAN) 17 December 1998 (1998-12-17)	1,49,52, 53
A	EP 0 254 241 A (YOSHITOMI PHARMACEUTICAL) 27 January 1988 (1988-01-27) claims 1,4	1,49,52, 53
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">20 March 2002</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">05/04/2002</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Alfaro Faus, I</div>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/25180

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9856785	A	17-12-1998	AU WO	7553198 A 9856785 A1
				30-12-1998 17-12-1998
EP 0254241	A	27-01-1988	EP JP JP JP US	0254241 A1 1904705 C 6015542 B 63146880 A 4808620 A
				27-01-1988 08-02-1995 02-03-1994 18-06-1988 28-02-1989